



## IGFL3 Polyclonal Antibody

play critical roles in cellular energy metabolism and in growth and development.		
Reactivity Human;Rat;Mouse;  Applications WB;ELISA  Gene Name IGFL3 UNQ483/PRO982  Protein Name Insulin growth factor-like family member 3  Immunogen Synthesized peptide derived from human protein . at AA range: 10-90  Specificity IGFL3 Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 13kD  Cell Pathway Secreted .  Tissue Specificity Detected in the cerebellum.  Function similarity-Belongs to the IGFL family.,  Background IGF like family member 3(IGFL3) Homo sapiens IGFL3 belongs to the insulin-like growth factor (IGF: see MIM 14/440) family of signaling molecules tha play critical roles in cellular energy metabolism and in growth and development especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]) [supplied by OhllM, Mar 2008],	Catalog No	BYab-05101
Applications WB;ELISA  Gene Name IGFL3 UNQ483/PRO982  Protein Name Insulin growth factor-like family member 3  Immunogen Synthesized peptide derived from human protein . at AA range: 10-90  Specificity IGFL3 Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 13kD  Cell Pathway Secreted .  Tissue Specificity Detected in the cerebellum.  Function similarity:Belongs to the IGFL family.,  Background IGF like family member 3(IGFL3) Homo sapiens IGFL3 belongs to the insulin-like growth factor (IGF; see MIM 147/440) family of signaling molecules tha play critical roles in cellular energy metabolism and in growth and development, especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]), [supplied by OMIM, Mar 2008],  matters needing Avoid repeated freezing and thawing!	Isotype	IgG
Gene Name IGFL3 UNQ483/PRO982 Protein Name Insulin growth factor-like family member 3 Immunogen Synthesized peptide derived from human protein . at AA range: 10-90 Specificity IGFL3 Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms  Observed Band 13kD  Cell Pathway Secreted .  Tissue Specificity Detected in the cerebellum.  Function similarity:Belongs to the IGFL family.,  Background IGF like family member 3(IGFL3) Homo sapiens IGFL3 belongs to the insulin-like growth factor (IGF; see MIM 147/440) family of signaling molecules tha play critical roles in cellular energy metabolism and in growth and development, especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]).[supplied by OMIM, Mar 2008],  matters needing Avoid repeated freezing and thawing!	Reactivity	Human;Rat;Mouse;
Protein Name         Insulin growth factor-like family member 3           Immunogen         Synthesized peptide derived from human protein . at AA range: 10-90           Specificity         IGFL3 Polyclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Polyclonal, Rabbit, IgG           Purification         The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000 ELISA 1:5000-20000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         13kD           Cell Pathway         Secreted .           Tissue Specificity         Detected in the cerebellum.           Function         similarity:Belongs to the IGFL family.,           Background         IGF like family member 3(IGFL3) Homo sapiens         IGFL3 belongs to the insulin-like growth factor (IGF; see MIM 147440) family of signaling molecules tha play critical roles in cellular energy metabolism and in growth and development, especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]). [supplied by OMIM, Mar 2008],           matters needing         Avoid repeated freezing and thawing!	Applications	WB;ELISA
Immunogen Synthesized peptide derived from human protein . at AA range: 10-90  Specificity IGFL3 Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 13kD  Cell Pathway Secreted .  Tissue Specificity Detected in the cerebellum.  Function similarity:Belongs to the IGFL family.,  Background IGF like family member 3(IGFL3) Homo sapiens IGFL3 belongs to the insulin-like growth factor (IGF; see MIM 147/440) family signaling molecules tha play critical roles in cellular energy metabolism and in growth and development, especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]).[supplied by OMIM, Mar 2008],  matters needing Avoid repeated freezing and thawing!	Gene Name	IGFL3 UNQ483/PRO982
Specificity       IGFL3 Polyclonal Antibody detects endogenous levels of protein.         Formulation       Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.         Source       Polyclonal, Rabbit, IgG         Purification       The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000 ELISA 1:5000-20000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band       13kD         Cell Pathway       Secreted .         Tissue Specificity       Detected in the cerebellum.         Function       similarity:Belongs to the IGFL family.,         Background       IGF like family member 3(IGFL3) Homo sapiens IGFL3 belongs to the insulin-like growth factor (IGF; see MIM 147440) family of signaling molecules tha play critical roles in cellular energy metabolism and in growth and development, especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]), [supplied by OMIM, Mar 2008],         matters needing       Avoid repeated freezing and thawing!	Protein Name	Insulin growth factor-like family member 3
Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source  Polyclonal, Rabbit, IgG  Purification  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000 ELISA 1:5000-20000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  13kD  Cell Pathway  Secreted.  Tissue Specificity  Detected in the cerebellum.  Function  similarity:Belongs to the IGFL family.,  Background  IGF like family member 3(IGFL3) Homo sapiens IGFL3 belongs to the insulin-like growth factor (IGF; see MIM 147440) family of signaling molecules tha play critical roles in cellular energy metabolism and in growth and development, especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]).[supplied by OMIM, Mar 2008],  matters needing  Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from human protein . at AA range: 10-90
Source       Polyclonal, Rabbit,IgG         Purification       The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000 ELISA 1:5000-20000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band       13kD         Cell Pathway       Secreted .         Tissue Specificity       Detected in the cerebellum.         Function       similarity:Belongs to the IGFL family.,         Background       IGF like family member 3(IGFL3) Homo sapiens IGFL3 belongs to the insulin-like growth factor (IGF; see MIM 147440) family of signaling molecules tha play critical roles in cellular energy metabolism and in growth and development, especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]).[supplied by OMIM, Mar 2008],         matters needing       Avoid repeated freezing and thawing!	Specificity	IGFL3 Polyclonal Antibody detects endogenous levels of protein.
Purification  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000 ELISA 1:5000-20000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  13kD  Cell Pathway  Secreted .  Tissue Specificity  Detected in the cerebellum.  Function  similarity:Belongs to the IGFL family.,  Background  IGF like family member 3(IGFL3) Homo sapiens IGFL3 belongs to the insulin-like growth factor (IGF; see MIM 147/440) family of signaling molecules tha play critical roles in cellular energy metabolism and in growth and development, especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]).[supplied by OMIM, Mar 2008],  matters needing  Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 13kD  Cell Pathway Secreted.  Tissue Specificity Detected in the cerebellum.  Function similarity:Belongs to the IGFL family.,  Background IGF like family member 3(IGFL3) Homo sapiens IGFL3 belongs to the insulin-like growth factor (IGF; see MIM 147440) family of signaling molecules that play critical roles in cellular energy metabolism and in growth and development, especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]).[supplied by OMIM, Mar 2008],  matters needing Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 13kD  Cell Pathway Secreted.  Tissue Specificity Detected in the cerebellum.  Function similarity:Belongs to the IGFL family.,  Background IGF like family member 3(IGFL3) Homo sapiens IGFL3 belongs to the insulin-like growth factor (IGF; see MIM 147440) family of signaling molecules that play critical roles in cellular energy metabolism and in growth and development, especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]).[supplied by OMIM, Mar 2008],  matters needing Avoid repeated freezing and thawing!	Purification	•
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 13kD  Cell Pathway Secreted .  Tissue Specificity Detected in the cerebellum.  Function similarity:Belongs to the IGFL family.,  Background IGF like family member 3(IGFL3) Homo sapiens IGFL3 belongs to the insulin-like growth factor (IGF; see MIM 147440) family of signaling molecules that play critical roles in cellular energy metabolism and in growth and development, especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]).[supplied by OMIM, Mar 2008],  matters needing Avoid repeated freezing and thawing!	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability  Synonyms  Observed Band  13kD  Cell Pathway  Secreted.  Tissue Specificity  Detected in the cerebellum.  Function  Similarity:Belongs to the IGFL family.,  Background  IGF like family member 3(IGFL3) Homo sapiens IGFL3 belongs to the insulin-like growth factor (IGF; see MIM 147440) family of signaling molecules tha play critical roles in cellular energy metabolism and in growth and development, especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]).[supplied by OMIM, Mar 2008],  matters needing  Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms  Observed Band 13kD  Cell Pathway Secreted.  Tissue Specificity Detected in the cerebellum.  Function similarity:Belongs to the IGFL family.,  Background IGF like family member 3(IGFL3) Homo sapiens IGFL3 belongs to the insulin-like growth factor (IGF; see MIM 147440) family of signaling molecules that play critical roles in cellular energy metabolism and in growth and development, especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]).[supplied by OMIM, Mar 2008],  matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band  Cell Pathway  Secreted .  Tissue Specificity  Detected in the cerebellum.  Function  Similarity:Belongs to the IGFL family.,  Background  IGF like family member 3(IGFL3) Homo sapiens IGFL3 belongs to the insulin-like growth factor (IGF; see MIM 147440) family of signaling molecules that play critical roles in cellular energy metabolism and in growth and development, especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]).[supplied by OMIM, Mar 2008],  matters needing  Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway       Secreted .         Tissue Specificity       Detected in the cerebellum.         Function       similarity:Belongs to the IGFL family.,         Background       IGF like family member 3(IGFL3) Homo sapiens insulin-like growth factor (IGF; see MIM 147440) family of signaling molecules that play critical roles in cellular energy metabolism and in growth and development, especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]).[supplied by OMIM, Mar 2008],         matters needing       Avoid repeated freezing and thawing!	Synonyms	
Tissue Specificity  Detected in the cerebellum.  Function  Similarity:Belongs to the IGFL family.,  Background  IGF like family member 3(IGFL3) Homo sapiens IGFL3 belongs to the insulin-like growth factor (IGF; see MIM 147440) family of signaling molecules that play critical roles in cellular energy metabolism and in growth and development, especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]).[supplied by OMIM, Mar 2008],  matters needing  Avoid repeated freezing and thawing!	Observed Band	13kD
Function similarity:Belongs to the IGFL family.,  Background IGF like family member 3(IGFL3) Homo sapiens IGFL3 belongs to the insulin-like growth factor (IGF; see MIM 147440) family of signaling molecules that play critical roles in cellular energy metabolism and in growth and development, especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]).[supplied by OMIM, Mar 2008],  matters needing Avoid repeated freezing and thawing!	Cell Pathway	Secreted .
Background  IGF like family member 3(IGFL3) Homo sapiens IGFL3 belongs to the insulin-like growth factor (IGF; see MIM 147440) family of signaling molecules that play critical roles in cellular energy metabolism and in growth and development, especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]).[supplied by OMIM, Mar 2008],  matters needing  Avoid repeated freezing and thawing!	Tissue Specificity	Detected in the cerebellum.
insulin-like growth factor (IGF; see MIM 147440) family of signaling molecules that play critical roles in cellular energy metabolism and in growth and development, especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]).[supplied by OMIM, Mar 2008],  matters needing  Avoid repeated freezing and thawing!	Function	similarity:Belongs to the IGFL family.,
	Background	insulin-like growth factor (IGF; sée MIM 147440) family of signaling molecules that play critical roles in cellular energy metabolism and in growth and development, especially prenatal growth (Emtage et al., 2006 [PubMed 16890402]).[supplied by
	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

