



PDCD7 Polyclonal Antibody

Catalog NoBYab-05148IsotypeIgGReactivityHuman;MouseApplicationsWB;ELISAGene NamePDCD7Protein NameProgrammed cell death protein 7 (ES18) (hES18)ImmunogenSynthesized peptide derived from human protein . at AA range: 300-380SpecificityPDCD7 Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/mlPurity≥90%
Reactivity Human;Mouse Applications WB;ELISA Gene Name PDCD7 Protein Name Programmed cell death protein 7 (ES18) (hES18) Immunogen Synthesized peptide derived from human protein . at AA range: 300-380 Specificity PDCD7 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml
Applications WB;ELISA Gene Name PDCD7 Protein Name Programmed cell death protein 7 (ES18) (hES18) Immunogen Synthesized peptide derived from human protein . at AA range: 300-380 Specificity PDCD7 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,lgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml
Gene NamePDCD7Protein NameProgrammed cell death protein 7 (ES18) (hES18)ImmunogenSynthesized peptide derived from human protein . at AA range: 300-380SpecificityPDCD7 Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit, lgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/ml
Protein NameProgrammed cell death protein 7 (ES18) (hES18)ImmunogenSynthesized peptide derived from human protein . at AA range: 300-380SpecificityPDCD7 Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit, lgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/ml
ImmunogenSynthesized peptide derived from human protein . at AA range: 300-380SpecificityPDCD7 Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/ml
SpecificityPDCD7 Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/ml
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml
Concentration 1 mg/ml
Purity ≥90%
Storage Stability -20°C/1 year
Synonyms
Observed Band 53kD
Cell Pathway Nucleus .
Tissue Specificity Duodenum, Placenta, Small intestine, Thymus,
Function function:Promotes apoptosis when overexpressed.,subunit:Interacts with RBM40.,
This gene encodes a 59 kDa protein that is associated with the U11 small nuclear ribonucleoprotein (snRNP), which is a component of the minor U12-typ spliceosome responsible for catalyzing pre-mRNA splicing of U12-type introns. [provided by RefSeq, Dec 2010],
matters needing Avoid repeated freezing and thawing! attention

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



ı	Isaq	P S	ua	ae	stic	าทร
u	Jau	C 3	uч	uc.	วแง	ノロコ

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images	

Nanjing BYabscience technology Co.,Ltd