



## NRX1B Polyclonal Antibody

Catalog No	BYab-05824
Isotype	lgG
Reactivity	Human;Rat;Mouse
Applications	WB;ELISA
Gene Name	NRXN1
Protein Name	Neurexin-1-beta (Neurexin I-beta)
Immunogen	Synthesized peptide derived from human protein . at AA range: 260-340
Specificity	NRX1B Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	48kD
Cell Pathway	Cell junction, synapse, presynaptic cell membrane ; Single-pass type I membrane protein .
Tissue Specificity	
Function	alternative products: A number of isoforms are produced by alternative promoter usage including the alpha-type (shown here) and beta-type (AC P58400) isoforms which differ in their N-terminus. Additional isoforms may be produced by alternative splicing, alternative products: A number of isoforms, alpha-type (AC Q9ULB1) and beta-type (shown here), are produced by alternative promoter usage. Beta-type isoforms differ from alpha-type isoforms in their N-terminus, function: Neuronal cell surface protein that may be involved in cell recognition and cell adhesion by forming intracellular junctions through binding to neuroligins. May play a role in formation or maintenance of synaptic junctions. May mediate intracellular signaling., function: Neuronal cell surface protein that may be involved in cell recognition and cell adhesion. May mediate intracellular signaling., PTM: Highly O-glycosylated and mino
	Newing Dyahasianas tashnalamu Calltd

## Nanjing BYabscience technology Co.,Ltd

网址:www.njbybio.com 官方热线:025-5229-8998 监督电话:15950492658

<b>博研生物</b> BYabscience
----------------------------



Background	alternative products: A number of isoforms are produced by alternative promoter usage including the alpha-type (shown here) and beta-type (AC P58400) isoforms which differ in their N-terminus. Additional isoforms may be produced by alternative splicing, alternative products: A number of isoforms, alpha-type (AC Q9ULB1) and beta-type (shown here), are produced by alternative promoter usage. Beta-type isoforms differ from alpha-type isoforms in their N-terminus, function: Neuronal cell surface protein that may be involved in cell recognition and cell adhesion by forming intracellular junctions through binding to neuroligins. May play a role in formation or maintenance of synaptic junctions. May mediate intracellular signaling., function: Neuronal cell surface protein that may be involved in cell recognition and cell adhesion. May mediate intracellular signaling., PTM:Highly O-glycosylated and minor N-glycosylated., PTM:N- and O-glycosylated., similarity: Belongs to the neurexin family., similarity: Contains 1 laminin G-like domain., similarity: Contains 3 EGF-like domains., similarity: Contains 6 laminin G-like domains., subunit: The cytoplasmic C-terminal region binds to CASK, CASKIN1 and APBA1. The laminin G-like domain 2 binds to NXPH1. Specific isoforms bind to alpha-dystroglycan and to alpha-latroxin. Interacts with SYT13 and SYTL1., subunit: The cytoplasmic C-terminal region binds to CASK. Isoforms Beta 4b bind neuroligins NLGN1, NLGN2 and NLGN3, alpha-dystroglycan and alpha-latrotoxin., tissue specificity:Heart and brain.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

Nanjing BYabscience technology Co.,Ltd