



ZDH13 Polyclonal Antibody

Catalog No	BYab-06693
Isotype	lgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	ZDHHC13 HIP14L HIP3RP
Protein Name	Palmitoyltransferase ZDHHC13 (EC 2.3.1) (Huntingtin-interacting protein 14-related protein) (HIP14-related protein) (Huntingtin-interacting protein HIP3RP) (Putative MAPK-activating protein PM03) (P
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	ZDH13 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	
-	≥90%
Storage Stability	≥90%
Storage Stability Synonyms	≥90% -20°C/1 year
Storage Stability Synonyms Observed Band	≥90% -20°C/1 year 68kD Golgi apparatus membrane ; Multi-pass membrane protein . Cytoplasmic vesicle membrane ; Multi-pass membrane protein . Low extracellular Mg(2+) induces
Storage Stability Synonyms Observed Band Cell Pathway	 ≥90% -20°C/1 year 68kD Golgi apparatus membrane ; Multi-pass membrane protein . Cytoplasmic vesicle membrane ; Multi-pass membrane protein . Low extracellular Mg(2+) induces increase in Golgi and in post-Golgi vesicles Eye,Lung fibroblast,Placenta,Teratocarcinoma,Testis, catalytic activity:Palmitoyl-CoA + protein-cysteine = S-palmitoyl protein + CoA.,domain:The DHHC domain is required for palmitoyltransferase
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	 ≥90% -20°C/1 year 68kD Golgi apparatus membrane ; Multi-pass membrane protein . Cytoplasmic vesicle membrane ; Multi-pass membrane protein . Low extracellular Mg(2+) induces increase in Golgi and in post-Golgi vesicles Eye,Lung fibroblast,Placenta,Teratocarcinoma,Testis, catalytic activity:Palmitoyl-CoA + protein-cysteine = S-palmitoyl protein + CoA.,domain:The DHHC domain is required for palmitoyltransferase activity.,function:May be involved in the NF-kappa-B signaling pathway.,sequence caution:Translated as Cys.,similarity:Belongs to the DHHC palmitoyltransferase family. AKR/ZDHHC17 subfamily.,similarity:Contains 1 DHHC-type zinc



国内优质抗体供应商

Шī, 精准的 WB 检测服务

24H 在线服务,欢迎咨询

	caution:Translated as Cys.,similarity:Belongs to the DHHC palmitoyltransferase family. AKR/ZDHHC17 subfamily.,similarity:Contains 1 DHHC-type zinc finger.,similarity:Contains 6 ANK repeats.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd