



Unc18-1 Polyclonal Antibody

| BYab-12835 |
|--|
| lgG |
| Human;Mouse;Rat |
| IHC;IF;ELISA |
| STXBP1 |
| Syntaxin-binding protein 1 |
| The antiserum was produced against synthesized peptide derived from human MUNC-18a. AA range:279-328 |
| Unc18-1 Polyclonal Antibody detects endogenous levels of Unc18-1 protein. |
| Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Polyclonal, Rabbit,IgG |
| The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications. |
| 1 mg/ml |
| ≥90% |
| -20°C/1 year |
| STXBP1; UNC18A; Syntaxin-binding protein 1; MUNC18-1; N-Sec1; Protein unc-18 homolog 1; Unc18-1; Protein unc-18 homolog A; Unc-18A; p67 |
| |
| Cytoplasm, cytosol . Membrane; Peripheral membrane protein. |
| Brain and spinal cord. Highly enriched in axons. |
| disease:Defects in STXBP1 are the cause of early infantile epileptic encephalopathy type 4 (EIEE4) [MIM:612164]. Affected individuals have neonatal or infantile onset of seizures, suppression-burst pattern on EEG, profound mental retardation, and MRI evidence of hypomyelination.,function:May participate in the regulation of synaptic vesicle docking and fusion, possibly through interaction with GTP-binding proteins. Essential for neurotransmission and binds syntaxin, a component of the synaptic vesicle fusion machinery probably in a 1:1 ratio. Can interact with syntaxins 1, 2, and 3 but not syntaxin 4. May play a role in determining the specificity of intracellular fusion reactions.,similarity:Belongs to |
| |

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658






| Background | This gene encodes a syntaxin-binding protein. The encoded protein appears to play a role in release of neurotransmitters via regulation of syntaxin, a transmembrane attachment protein receptor. Mutations in this gene have been associated with infantile epileptic encephalopathy-4. Alternatively spliced transcript variants have been described. [provided by RefSeq, Feb 2010], |
|---------------------------|---|
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |

specificity: Brain and spinal cord. Highly enriched in axons...

Products Images



Immunofluorescence analysis of NIH/3T3 cells, using MUNC-18a Antibody. The picture on the right is blocked with the synthesized peptide.

Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using MUNC-18a Antibody. The picture on the right is blocked with the synthesized peptide.

Nanjing BYabscience technology Co.,Ltd