



## FAM3C Polyclonal Antibody

Catalog No         BYab-06869           Isotype         IgG           Reactivity         Human;Mouse;Rat           Applications         WB;ELISA           Gene Name         FAM3C ILEI GS3786           Protein Name         Protein FAM3C (Interleukin-like EMT inducer)           Immunogen         Synthesized peptide derived from part region of human protein           Specificity         FAM3C Polyclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Polyclonal, Rabbit, IgG           Purification         The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000 ELISA 1:5000-20000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Synonyms           Observed Band         24kD           Cell Pathway         Secreted . Cytoplasmic vesicle . Cytoplasmic in some cancer cells.           Tissue Specificity         Present in most secretory epithelia (at protein level).           Function         similarity:Belongs to the FAM3 family., tissue specificity:Ubiquitous.,           Background		
Reactivity Human;Mouse;Rat  Applications WB;ELISA  Gene Name FAM3C ILEI GS3786 Protein Name Protein FAM3C (Interleukin-like EMT inducer) Immunogen Synthesized peptide derived from part region of human protein  Specificity FAM3C Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms  Observed Band 24kD  Cell Pathway Secreted . Cytoplasmic vesicle . Cytoplasmic in some cancer cells.  Tissue Specificity Present in most secretory epithelia (at protein level).  Function similarity:Belongs to the FAM3 family, tissue specificity:Ubiquitous.,  This gene is a member of the family with sequence similarity 3 (FAM3) family and encodes a secreted protein with a GG domain. A change in expression of this protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq, Mar 2010],  matters needing Avoid repeated freezing and thawing!	Catalog No	BYab-06869
Applications WB;ELISA  Gene Name FAM3C ILEI GS3786  Protein Name Protein FAM3C (Interleukin-like EMT inducer)  Immunogen Synthesized peptide derived from part region of human protein  Specificity FAM3C Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 24kD  Cell Pathway Secreted . Cytoplasmic vesicle . Cytoplasmic in some cancer cells.  Tissue Specificity Present in most secretory epithelia (at protein level).  Function similarity:Belongs to the FAM3 family.,tissue specificity:Ubiquitous.,  This gene is a member of the family with sequence similarity 3 (FAM3) family and encodes a secreted protein with a GG domain. A change in expression of this protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq, Mar 2010],  matters needing Avoid repeated freezing and thawing!	Isotype	IgG
Gene Name FAM3C ILEI GS3786  Protein Name Protein FAM3C (Interleukin-like EMT inducer)  Immunogen Synthesized peptide derived from part region of human protein  Specificity FAM3C Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 24kD  Cell Pathway Secreted . Cytoplasmic vesicle . Cytoplasmic in some cancer cells.  Tissue Specificity Present in most secretory epithelia (at protein level).  Function similarity:Belongs to the FAM3 family.,tissue specificity:Ubiquitous.,  This gene is a member of the family with sequence similarity 3 (FAM3) family and encodes a secreted protein with a GG domain. A change in expression of this protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq, Mar 2010],  matters needing Avoid repeated freezing and thawing!	Reactivity	Human;Mouse;Rat
Protein Name         Protein FAM3C (Interleukin-like EMT inducer)           Immunogen         Synthesized peptide derived from part region of human protein           Specificity         FAM3C Polyclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Polyclonal, Rabbit, IgG           Purification         The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000 ELISA 1:5000-20000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         24kD           Cell Pathway         Secreted . Cytoplasmic vesicle . Cytoplasmic in some cancer cells.           Tissue Specificity         Present in most secretory epithelia (at protein level).           Function         similarity:Belongs to the FAM3 family.,tissue specificity:Ubiquitous.,           Background         This gene is a member of the family with sequence similarity 3 (FAM3) family and encodes a secreted protein with a GG domain. A change in expression of this protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq, Mar 2010],           matters needing         Avoid repeated freezing and thawing!	Applications	WB;ELISA
Immunogen       Synthesized peptide derived from part region of human protein         Specificity       FAM3C Polyclonal Antibody detects endogenous levels of protein.         Formulation       Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.         Source       Polyclonal, Rabbit, IgG         Purification       The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000 ELISA 1:5000-20000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band       24kD         Cell Pathway       Secreted . Cytoplasmic vesicle . Cytoplasmic in some cancer cells.         Tissue Specificity       Present in most secretory epithelia (at protein level).         Function       similarity:Belongs to the FAM3 family.,tissue specificity:Ubiquitous.,         Background       This gene is a member of the family with sequence similarity 3 (FAM3) family and encodes a secreted protein with a GG domain. A change in expression of this protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq, Mar 2010],         matters needing       Avoid repeated freezing and thawing!	Gene Name	FAM3C ILEI GS3786
Specificity FAM3C Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms  Observed Band 24kD  Cell Pathway Secreted . Cytoplasmic vesicle . Cytoplasmic in some cancer cells.  Tissue Specificity Present in most secretory epithelia (at protein level).  Function similarity:Belongs to the FAM3 family.,tissue specificity:Ubiquitous.,  Background This gene is a member of the family with sequence similarity 3 (FAM3) family and encodes a secreted protein with a GG domain. A change in expression of this protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq, Mar 2010], matters needing Avoid repeated freezing and thawing!	Protein Name	Protein FAM3C (Interleukin-like EMT inducer)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity 290% Storage Stability -20°C/1 year  Synonyms  Observed Band 24kD  Cell Pathway Secreted . Cytoplasmic vesicle . Cytoplasmic in some cancer cells.  Tissue Specificity Present in most secretory epithelia (at protein level).  Function similarity:Belongs to the FAM3 family, tissue specificity:Ubiquitous.,  Background This gene is a member of the family with sequence similarity 3 (FAM3) family and encodes a secreted protein with a GG domain. A change in expression of this protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq, Mar 2010],  matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from part region of human protein
Source       Polyclonal, Rabbit,IgG         Purification       The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000 ELISA 1:5000-20000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band       24kD         Cell Pathway       Secreted . Cytoplasmic vesicle . Cytoplasmic in some cancer cells.         Tissue Specificity       Present in most secretory epithelia (at protein level).         Function       similarity:Belongs to the FAM3 family.,tissue specificity:Ubiquitous.,         Background       This gene is a member of the family with sequence similarity 3 (FAM3) family and encodes a secreted protein with a GG domain. A change in expression of this protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq, Mar 2010],         matters needing       Avoid repeated freezing and thawing!	Specificity	FAM3C Polyclonal Antibody detects endogenous levels of protein.
Purification  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000 ELISA 1:5000-20000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  24kD  Cell Pathway  Secreted . Cytoplasmic vesicle . Cytoplasmic in some cancer cells.  Tissue Specificity  Present in most secretory epithelia (at protein level).  Function  similarity:Belongs to the FAM3 family.,tissue specificity:Ubiquitous.,  Background  This gene is a member of the family with sequence similarity 3 (FAM3) family and encodes a secreted protein with a GG domain. A change in expression of this protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq, Mar 2010],  matters needing  Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 24kD  Cell Pathway Secreted . Cytoplasmic vesicle . Cytoplasmic in some cancer cells.  Tissue Specificity Present in most secretory epithelia (at protein level).  Function similarity:Belongs to the FAM3 family.,tissue specificity:Ubiquitous.,  Background This gene is a member of the family with sequence similarity 3 (FAM3) family and encodes a secreted protein with a GG domain. A change in expression of this protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq, Mar 2010],  matters needing Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 24kD  Cell Pathway Secreted . Cytoplasmic vesicle . Cytoplasmic in some cancer cells.  Tissue Specificity Present in most secretory epithelia (at protein level).  Function similarity:Belongs to the FAM3 family.,tissue specificity:Ubiquitous.,  Background This gene is a member of the family with sequence similarity 3 (FAM3) family and encodes a secreted protein with a GG domain. A change in expression of this protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq, Mar 2010],  matters needing Avoid repeated freezing and thawing!	Purification	
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 24kD  Cell Pathway Secreted . Cytoplasmic vesicle . Cytoplasmic in some cancer cells.  Tissue Specificity Present in most secretory epithelia (at protein level).  Function similarity:Belongs to the FAM3 family.,tissue specificity:Ubiquitous.,  Background This gene is a member of the family with sequence similarity 3 (FAM3) family and encodes a secreted protein with a GG domain. A change in expression of this protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq, Mar 2010],  matters needing Avoid repeated freezing and thawing!	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability -20°C/1 year  Synonyms  Observed Band 24kD  Cell Pathway Secreted . Cytoplasmic vesicle . Cytoplasmic in some cancer cells.  Tissue Specificity Present in most secretory epithelia (at protein level).  Function similarity:Belongs to the FAM3 family.,tissue specificity:Ubiquitous.,  Background This gene is a member of the family with sequence similarity 3 (FAM3) family and encodes a secreted protein with a GG domain. A change in expression of this protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq, Mar 2010],  matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms  Observed Band 24kD  Cell Pathway Secreted . Cytoplasmic vesicle . Cytoplasmic in some cancer cells.  Tissue Specificity Present in most secretory epithelia (at protein level).  Function similarity:Belongs to the FAM3 family.,tissue specificity:Ubiquitous.,  Background This gene is a member of the family with sequence similarity 3 (FAM3) family and encodes a secreted protein with a GG domain. A change in expression of this protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq, Mar 2010],  matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band  Cell Pathway  Secreted . Cytoplasmic vesicle . Cytoplasmic in some cancer cells.  Tissue Specificity  Present in most secretory epithelia (at protein level).  Function  similarity:Belongs to the FAM3 family.,tissue specificity:Ubiquitous.,  This gene is a member of the family with sequence similarity 3 (FAM3) family and encodes a secreted protein with a GG domain. A change in expression of this protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq, Mar 2010],  matters needing  Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway       Secreted . Cytoplasmic vesicle . Cytoplasmic in some cancer cells.         Tissue Specificity       Present in most secretory epithelia (at protein level).         Function       similarity:Belongs to the FAM3 family.,tissue specificity:Ubiquitous.,         Background       This gene is a member of the family with sequence similarity 3 (FAM3) family and encodes a secreted protein with a GG domain. A change in expression of this protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq, Mar 2010],         matters needing       Avoid repeated freezing and thawing!	Synonyms	
Tissue Specificity Present in most secretory epithelia (at protein level).  Function similarity:Belongs to the FAM3 family.,tissue specificity:Ubiquitous.,  This gene is a member of the family with sequence similarity 3 (FAM3) family and encodes a secreted protein with a GG domain. A change in expression of this protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq, Mar 2010],  Mar 2010],  Avoid repeated freezing and thawing!	Observed Band	24kD
Function similarity:Belongs to the FAM3 family.,tissue specificity:Ubiquitous.,  Background This gene is a member of the family with sequence similarity 3 (FAM3) family and encodes a secreted protein with a GG domain. A change in expression of this protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq, Mar 2010],  matters needing Avoid repeated freezing and thawing!	Cell Pathway	Secreted . Cytoplasmic vesicle . Cytoplasmic in some cancer cells.
Background  This gene is a member of the family with sequence similarity 3 (FAM3) family and encodes a secreted protein with a GG domain. A change in expression of this protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq, Mar 2010],  matters needing  Avoid repeated freezing and thawing!	Tissue Specificity	Present in most secretory epithelia (at protein level).
protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq, Mar 2010],  matters needing Avoid repeated freezing and thawing!	Function	similarity:Belongs to the FAM3 family.,tissue specificity:Ubiquitous.,
	Background	protein has been noted in pancreatic cancer-derived cells. [provided by RefSeq,
	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658