



NEST Polyclonal Antibody

Catalog No	BYab-06913
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB;IHC
Gene Name	NES Nbla00170
Protein Name	Nestin
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	NEST Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year 178kD cytoplasm,intermediate filament,extracellular matrix,intermediate filament
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year 178kD cytoplasm,intermediate filament,extracellular matrix,intermediate filament cytoskeleton, CNS stem cells. developmental stage:Upon terminal neural differentiation, nestin is down-regulated and replaced by neurofilaments.,function:May play a role in the trafficking and distribution of IF proteins and potentially other cellular factors to daughter cells during progenitor cell division.,similarity:Belongs to the
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year 178kD cytoplasm,intermediate filament,extracellular matrix,intermediate filament cytoskeleton, CNS stem cells. developmental stage:Upon terminal neural differentiation, nestin is down-regulated and replaced by neurofilaments.,function:May play a role in the trafficking and distribution of IF proteins and potentially other cellular factors to daughter cells during progenitor cell division.,similarity:Belongs to the intermediate filament family.,subunit:Interacts with FHOD3.,tissue specificity:CNS
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year 178kD cytoplasm,intermediate filament,extracellular matrix,intermediate filament cytoskeleton, CNS stem cells. developmental stage:Upon terminal neural differentiation, nestin is down-regulated and replaced by neurofilaments.,function:May play a role in the trafficking and distribution of IF proteins and potentially other cellular factors to daughter cells during progenitor cell division.,similarity:Belongs to the intermediate filament family.,subunit:Interacts with FHOD3.,tissue specificity:CNS stem cells., This gene encodes a member of the intermediate filament protein family and is

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Mechanism of Pulp Regeneration Based on Concentrated Growth Factors Regulating Cell Differentiation Bioengineering-Basel Sijing Yu, Yi Zheng, Qiang Guo, Wenxu Li, Ling Ye, Bo Gao IHC Mouse human dental pulp cells (hDPCs)-Xenograft

Nanjing BYabscience technology Co.,Ltd