



## **TICN1** Polyclonal Antibody

Catalog NoBYab-07132IsotypeIgGReactivityHuman;MouseApplicationsWB;ELISAGene NameSPOCK1 SPOCK TIC1 TICN1Protein NameTestican-1 (Protein SPOCK)ImmunogenSynthesized peptide derived from human protein . at AA range: 350-430SpecificityTICN1 Polycional Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolycional, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:500-2000Concentration1 mg/mlPurity290%Storage Stability-20°C/1 yearStorage StabilitySecreted, extracellular space, extracellular matrix.Tissue SpecificityBrain, Cerebellum, Hippocampus, Seminal plasmaFunctionfunction: May play a role in cell-cell and cell-matrix interactions. May contribute to various neuronal mechanisms in the central nervous system., PTM:Contains chondrollin sulfate and heparan-sulfate Chinked omain., similarity:Contains 1 thyroglobulin type-1 domain.BackgroundThis gene encodes the protein core of a seminal plasma proteoglycan containing rhondrollin and heparan-sulfate chains. The protein&apors's function is unknown, atthough similarity to tropine protein core of a seminal plasma proteoglycan containing rhondrollin and heparan-sulfate chains. The protein&apors's function is unknown, atthough similarity to tropine. Face-steine protease-inhibitors sugges ts ite.		
ReactivityHuman;MouseApplicationsWB;ELISAGene NameSPOCK1 SPOCK TIC1 TICN1Protein NameTestican-1 (Protein SPOCK)ImmunogenSynthesized peptide derived from human protein . at AA range: 350-430SpecificityTICN1 Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsSecreted, extracellular space, extracellular matrix.Tissue SpecificityBrain, Cerebellum, Hippocampus, Seminal plasmaFunctionfunction:May play a role in cell-cell and cell-matrix interactions. May contribute to various neuronal mechanisms in the central nervous system, PTM:Contains chondrolitin- and heparan-sulfate ColinkedBackgroundThis gene encodes the protein core of a seminal plasma proteoglycan containing chondrolitin- and heparan-sulfate colains. The proteinsaport sungaes is in sunknown, atthough similarity to thropin-type cysteine proteinses-inhibitors suggests its sunknown,	Catalog No	BYab-07132
ApplicationsWB;ELISAGene NameSPOCK1 SPOCK TIC1 TICN1Protein NameTestican-1 (Protein SPOCK)ImmunogenSynthesized peptide derived from human protein . at AA range: 350-430SpecificityTICN1 Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsSecreted, extracellular space, extracellular matrix.Tissue SpecificityBrain,Cerebellum,Hippocampus,Seminal plasmaFunctionfunction:May play a role in cell-cell and cell-matrix interactions. May contribute to various neuronal mechanisms in the central nervous systemPTM:Contains chondroitin sulfate and heparan sulfate O-inked oligosaccharides.similarity:Contains 1 Kazal-like domainsimilarity:Contains 1 thyroglobulin type-1 domainBackgroundThis gene encodes the protein core of a seminal plasma proteoglycan containing chondroitin- and heparan-sulfate chains. The protease-inhibitors suggests its	Isotype	lgG
Gene NameSPOCK1 SPOCK TIC1 TICN1Protein NameTestican-1 (Protein SPOCK)ImmunogenSynthesized peptide derived from human protein . at AA range: 350-430SpecificityTICN1 Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCollege extracellular space, extracellular matrix.Tissue SpecificityBrain,Cerebellum,Hippocampus,Seminal plasmaFunctionfunction:May play a role in cell-cell and cell-matrix interactions. May contribute to various neuronal mechanisms in the central nervous system.,PTM:Contains chondroltin sulfate and heparan sulfate O-linked oligosaccharides.,similarity:Contains 1 Kazal-like domain.,similarity:Contains 1 thyroglobulin type-1 domain.	Reactivity	Human;Mouse
Protein NameTestican-1 (Protein SPOCK)ImmunogenSynthesized peptide derived from human protein . at AA range: 350-430SpecificityTICN1 Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsColserved Band48kDCell PathwaySecreted, extracellular space, extracellular matrix.Tissue SpecificityBrain,Cerebellum,Hippocampus,Seminal plasmaFunctionfunction:May play a role in cell-cell and cell-matrix interactions. May contribute to various neuronal mechanisms in the central nervous system.,PTM:Contains chondrollin sulfate and heparan sulfate C-linked oligosaccharides,similarity:Contains 1 Kazal-like domain.,similarity:Contains 1 thyroglobulin type-1 domain., athough similarity to thryopin-type cysteine protease-inhibitors suggests its	Applications	WB;ELISA
ImmunogenSynthesized peptide derived from human protein . at AA range: 350-430SpecificityTICN1 Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit, IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsSecreted, extracellular space, extracellular matrix.Tissue SpecificityBrain,Cerebellum,Hippocampus,Seminal plasmaFunctionUnction:May play a role in cell-cell and cell-matrix interactions. May contribute to various neuronal mechanisms in the cell-matrix interactions. May contribute to various neuronal mechanisms in the cell-matrix.FunctionUnction:May play a role in cell-cell and cell-matrix.BackgroundThis gene encodes the protein core of a seminal plasma proteoglycan containing chondroltin - and heparan-sulfate chains. The proteanse-inhibitors suggests its	Gene Name	SPOCK1 SPOCK TIC1 TICN1
Specificity       TICN1 Polyclonal Antibody detects endogenous levels of protein.         Formulation       Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.         Source       Polyclonal, Rabbit,IgG         Purification       The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000 ELISA 1:5000-20000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms	Protein Name	Testican-1 (Protein SPOCK)
FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCell PathwayCell PathwaySecreted, extracellular space, extracellular matrix.Tissue SpecificityBrain,Cerebellum,Hippocampus,Seminal plasmaFunctionfunction:May play a role in cell-cell and cell-matrix interactions. May contribute to various neuronal mechanisms in the central nervous system.,PTM:Contains 1 thyroglobulin type-1 domain.,BackgroundThis gene encodes the protein core of a seminal plasma proteoglycan containing chondroitin- and heparan-sulfate chains. The protein's function is unknown, although similarity to thyropin-type cysteine protease-inhibitors suggess is its	Immunogen	Synthesized peptide derived from human protein . at AA range: 350-430
SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsObserved Band48kDCell PathwaySecreted, extracellular space, extracellular matrix.Tissue SpecificityBrain,Cerebellum,Hippocampus,Seminal plasmaFunctionfunction:May play a role in cell-cell and cell-matrix interactions. May contribute to various neuronal mechanisms in the central nervous systemPTM:Contains 1 thyroglobulin type-1 domain.,BackgroundThis gene encodes the protein core of a seminal plasma proteoglycan containing chondroitin- and heparan-sulfate chains. The protein&apocis function is unknown, although similarity to thyropin-type cysteine protease-inhibitors suggests its	Specificity	TICN1 Polyclonal Antibody detects endogenous levels of protein.
Purification       The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000 ELISA 1:5000-20000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       -20°C/1 year         Observed Band       48kD         Cell Pathway       Secreted, extracellular space, extracellular matrix.         Tissue Specificity       Brain,Cerebellum,Hippocampus,Seminal plasma         Function       function:May play a role in cell-cell and cell-matrix interactions. May contribute to various neuronal mechanisms in the central nervous system.,PTM:Contains 1 thyroglobulin type-1 domain.,         Background       This gene encodes the protein core of a seminal plasma proteoglycan contains 1 thorgon synthetype cysteine protease-inhibitors suggests its	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000 ELISA 1:5000-20000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       -20°C/1 year         Observed Band       48kD         Cell Pathway       Secreted, extracellular space, extracellular matrix.         Tissue Specificity       Brain,Cerebellum,Hippocampus,Seminal plasma         Function       function:May play a role in cell-cell and cell-matrix interactions. May contribute to various neuronal mechanisms in the central nervous system.,PTM:Contains chondroitin sulfate and heparan sulfate O-linked oligosaccharides, similarity:Contains 1 Kazal-like domain.,similarity:Contains 1 thyroglobulin type-1 domain.,         Background       This gene encodes the protein core of a seminal plasma proteoglycan containing chondroitin- and heparan-sulfate chains. The protein's function is unknown, although similarity to thyropin-type cysteine protease-inhibitors suggests its	Source	Polyclonal, Rabbit,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       -20°C/1 year         Observed Band       48kD         Cell Pathway       Secreted, extracellular space, extracellular matrix.         Tissue Specificity       Brain,Cerebellum,Hippocampus,Seminal plasma         Function       function:May play a role in cell-cell and cell-matrix interactions. May contribute to various neuronal mechanisms in the central nervous system.,PTM:Contains chondroitin sulfate and heparan sulfate O-linked oligosaccharides.,similarity:Contains 1 Kazal-like domain.,similarity:Contains 1 thyroglobulin type-1 domain.,         Background       This gene encodes the protein core of a seminal plasma proteoglycan containing chondroitin- and heparan-sulfate chains. The protein's function is unknown, although similarity to thyropin-type cysteine protease-inhibitors suggests its	Purification	
Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       -20°C/1 year         Observed Band       48kD         Cell Pathway       Secreted, extracellular space, extracellular matrix.         Tissue Specificity       Brain,Cerebellum,Hippocampus,Seminal plasma         Function       function:May play a role in cell-cell and cell-matrix interactions. May contribute to various neuronal mechanisms in the central nervous system.,PTM:Contains chondroitin sulfate and heparan sulfate O-linked oligosaccharides.,similarity:Contains 1 Kazal-like domain.,similarity:Contains 1 thyroglobulin type-1 domain.,         Background       This gene encodes the protein core of a seminal plasma proteoglycan containing chondroitin- and heparan-sulfate chains. The protein's function is unknown, although similarity to thyropin-type cysteine protease-inhibitors suggests its	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability       -20°C/1 year         Synonyms       -20°C/1 year         Observed Band       48kD         Cell Pathway       Secreted, extracellular space, extracellular matrix.         Tissue Specificity       Brain,Cerebellum,Hippocampus,Seminal plasma         Function       function:May play a role in cell-cell and cell-matrix interactions. May contribute to various neuronal mechanisms in the central nervous system.,PTM:Contains chondroitin sulfate and heparan sulfate O-linked oligosaccharides.,similarity:Contains 1 Kazal-like domain.,similarity:Contains 1         Background       This gene encodes the protein core of a seminal plasma proteoglycan containing chondroitin- and heparan-sulfate chains. The protein's function is unknown, although similarity to thyropin-type cysteine protease-inhibitors suggests its	Concentration	1 mg/ml
Synonyms         Observed Band       48kD         Cell Pathway       Secreted, extracellular space, extracellular matrix.         Tissue Specificity       Brain,Cerebellum,Hippocampus,Seminal plasma         Function       function:May play a role in cell-cell and cell-matrix interactions. May contribute to various neuronal mechanisms in the central nervous systemPTM:Contains chondroitin sulfate and heparan sulfate O-linked oligosaccharidessimilarity:Contains 1 Kazal-like domainsimilarity:Contains 1 thyroglobulin type-1 domain.,         Background       This gene encodes the protein core of a seminal plasma proteoglycan containing chondroitin- and heparan-sulfate chains. The protein's function is unknown, although similarity to thyropin-type cysteine protease-inhibitors suggests its	Purity	≥90%
Observed Band48kDCell PathwaySecreted, extracellular space, extracellular matrix.Tissue SpecificityBrain,Cerebellum,Hippocampus,Seminal plasmaFunctionfunction:May play a role in cell-cell and cell-matrix interactions. May contribute to various neuronal mechanisms in the central nervous system.,PTM:Contains chondroitin sulfate and heparan sulfate O-linked oligosaccharides.,similarity:Contains 1 Kazal-like domain.,similarity:Contains 1 thyroglobulin type-1 domain.,BackgroundThis gene encodes the protein core of a seminal plasma proteoglycan containing chondroitin- and heparan-sulfate chains. The protein's function is unknown, although similarity to thyropin-type cysteine protease-inhibitors suggests its	Storage Stability	-20°C/1 year
Cell PathwaySecreted, extracellular space, extracellular matrix.Tissue SpecificityBrain,Cerebellum,Hippocampus,Seminal plasmaFunctionfunction:May play a role in cell-cell and cell-matrix interactions. May contribute to various neuronal mechanisms in the central nervous system.,PTM:Contains chondroitin sulfate and heparan sulfate O-linked oligosaccharides.,similarity:Contains 1 Kazal-like domain.,similarity:Contains 1 thyroglobulin type-1 domain.,BackgroundThis gene encodes the protein core of a seminal plasma proteoglycan containing chondroitin- and heparan-sulfate chains. The protein's function is unknown, although similarity to thyropin-type cysteine protease-inhibitors suggests its	Synonyms	
Tissue SpecificityBrain,Cerebellum,Hippocampus,Seminal plasmaFunctionfunction:May play a role in cell-cell and cell-matrix interactions. May contribute to various neuronal mechanisms in the central nervous system.,PTM:Contains chondroitin sulfate and heparan sulfate O-linked oligosaccharides.,similarity:Contains 1 Kazal-like domain.,similarity:Contains 1 thyroglobulin type-1 domain.,BackgroundThis gene encodes the protein core of a seminal plasma proteoglycan containing chondroitin- and heparan-sulfate chains. The protein's function is unknown, although similarity to thyropin-type cysteine protease-inhibitors suggests its	Observed Band	48kD
Functionfunction:May play a role in cell-cell and cell-matrix interactions. May contribute to various neuronal mechanisms in the central nervous system.,PTM:Contains chondroitin sulfate and heparan sulfate O-linked oligosaccharides.,similarity:Contains 1 Kazal-like domain.,similarity:Contains 1 thyroglobulin type-1 domain.,BackgroundThis gene encodes the protein core of a seminal plasma proteoglycan containing chondroitin- and heparan-sulfate chains. The protein's function is unknown, although similarity to thyropin-type cysteine protease-inhibitors suggests its	Cell Pathway	Secreted, extracellular space, extracellular matrix.
<ul> <li>various neuronal mechanisms in the central nervous system.,PTM:Contains chondroitin sulfate and heparan sulfate O-linked oligosaccharides.,similarity:Contains 1 Kazal-like domain.,similarity:Contains 1 thyroglobulin type-1 domain.,</li> <li>Background</li> <li>This gene encodes the protein core of a seminal plasma proteoglycan containing chondroitin- and heparan-sulfate chains. The protein's function is unknown, although similarity to thyropin-type cysteine protease-inhibitors suggests its</li> </ul>	Tissue Specificity	Brain,Cerebellum,Hippocampus,Seminal plasma
chondroitin- and heparan-sulfate chains. The protein's function is unknown, although similarity to thyropin-type cysteine protease-inhibitors suggests its	Function	various neuronal mechanisms in the central nervous system.,PTM:Contains chondroitin sulfate and heparan sulfate O-linked oligosaccharides.,similarity:Contains 1 Kazal-like domain.,similarity:Contains 1
	Background	chondroitin- and heparan-sulfate chains. The protein's function is unknown, although similarity to thyropin-type cysteine protease-inhibitors suggests its

Nanjing BYabscience technology Co.,Ltd





## matters needing attention

## Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images** 

Nanjing BYabscience technology Co.,Ltd