



## **OR1L8 Polyclonal Antibody**

Catalog No         BYab-07458           Isotype         IgG           Reactivity         Human;Rat;Mouse;           Applications         WB;ELISA           Gene Name         OR1L8           Protein Name         Olfactory receptor 1L8 (Olfactory receptor OR9-24)           Immunogen         Synthesized peptide derived from part region of human protein           Specificity         OR1L8 Polyclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Polyclonal, Rabbit, IgG           Purification         The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000 ELISA 1:5000-20000           Concentration         1 mg/ml           Purity         290%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         33kD           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Tissue Specificity         Function         function:Odorant receptor .;similarity:Belongs to the G-protein coupled familly.           Background         olfactory receptor family 1 subfamily L member 8(OR1L8) Homo sapien Olfactory receptors interact with odorant molecules in the nose, to inting neuronal respo	
Reactivity Human;Rat;Mouse;  Applications WB;ELISA  Gene Name OR1L8  Protein Name Olfactory receptor 1L8 (Olfactory receptor OR9-24)  Immunogen Synthesized peptide derived from part region of human protein  Specificity OR1L8 Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 33kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled ramily.,  Background olfactory receptor family 1 subfamily L member 8(OR1L8) Homo sapien olfactory receptors interact with odorant molecules in the nose, to initia neuronal response that triggers the perception of a smell. The olfactory proteins are members of a large family of-protein-coupled receptors of proteins are members of a large family of-protein-coupled receptors	
Applications WB;ELISA  Gene Name OR1L8  Protein Name Olfactory receptor 1L8 (Olfactory receptor OR9-24)  Immunogen Synthesized peptide derived from part region of human protein  Specificity OR1L8 Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 33kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled ramily.,  Background olfactory receptor family 1 subfamily L member 8(OR1L8) Homo sapien olfactory receptors interact with odorant molecules in the nose, to initia neuronal response that triggers the perception of a smell. The olfactory proteins are members of a large family of-protein-coupled receptors of the greeptors of a smell. The olfactory receptors are members of a large family of-protein-coupled receptors of the greeptors of the greeptors of a smell. The olfactory receptors are members of a large family of-protein-coupled receptors of the greeptors of the greeptor of the greeptors of the greeptor of the greeptors of the greeptors of the greeptor of t	
Gene Name         OR1L8           Protein Name         Olfactory receptor 1L8 (Olfactory receptor OR9-24)           Immunogen         Synthesized peptide derived from part region of human protein           Specificity         OR1L8 Polyclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Polyclonal, Rabbit, IgG           Purification         The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000 ELISA 1:5000-20000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         33kD           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Tissue Specificity         Function         function:Odorant receptor .,similarity:Belongs to the G-protein coupled refamily.,           Background         Olfactory receptor family 1 subfamily L member 8(OR1L8) Homo sapien Olfactory receptors interact with odorant molecules in the nose, to initia neuronal response that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors (specific secretors)	
Protein Name         Olfactory receptor 1L8 (Olfactory receptor OR9-24)           Immunogen         Synthesized peptide derived from part region of human protein           Specificity         OR1L8 Polyclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Polyclonal, Rabbit,IgG           Purification         The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000 ELISA 1:5000-20000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         33kD           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Tissue Specificity         Function           Function         function:Odorant receptor .,similarity:Belongs to the G-protein coupled family.           Background         Olfactory receptor family 1 subfamily L member 8(OR1L8) Homo sapier Olfactory receptors interact with odorant molecules in the nose, to initia neuronal response that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors of the perceptor of the protein-coupled receptors of the perceptor of the	
Immunogen         Synthesized peptide derived from part region of human protein           Specificity         OR1L8 Polyclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Polyclonal, Rabbit, IgG           Purification         The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000 ELISA 1:5000-20000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         33kD           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Tissue Specificity         Function         function:Odorant receptor .,similarity:Belongs to the G-protein coupled to family.,           Background         olfactory receptor family 1 subfamily L member 8(OR1L8) Homo sapier Olfactory receptors interact with odorant molecules in the nose, to initianeuronal response that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors of a large family of G-protein-coupled receptors in the coupled receptor of a large family of G-protein-coupled receptors in the coupled receptor of a large family of G-protein-coupled receptors in the coupled receptor of a large family of G-protein-coupled receptors in the coupled receptor of a large family of G-protein-coupled receptor of a large	
Specificity         OR1L8 Polyclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Polyclonal, Rabbit, IgG           Purification         The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000 ELISA 1:5000-20000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band           Observed Band         33kD           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Tissue Specificity         function:Odorant receptor .,similarity:Belongs to the G-protein coupled in family.,           Background         olfactory receptor family 1 subfamily L member 8(OR1L8) Homo sapien Olfactory receptors interact with odorant molecules in the nose, to initian neuronal response that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors (of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors (of protein-coupled receptors)	
Formulation       Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.         Source       Polyclonal, Rabbit, lgG         Purification       The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000 ELISA 1:5000-20000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band         Cell Pathway       Cell membrane; Multi-pass membrane protein.         Tissue Specificity       function:Odorant receptor .,similarity:Belongs to the G-protein coupled in family.,         Background       olfactory receptor family 1 subfamily L member 8(OR1L8) Homo sapien Olfactory receptors interact with odorant molecules in the nose, to initian neuronal response that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors (proteins are members of a large family of G-protein-coupled receptors (proteins are members of a large family of G-protein-coupled receptors (proteins are members of a large family of G-protein-coupled receptors (protein-coupled receptors (protein-coupled receptors)	
Source       Polyclonal, Rabbit, IgG         Purification       The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000 ELISA 1:5000-20000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band       33kD         Cell Pathway       Cell membrane; Multi-pass membrane protein.         Tissue Specificity       Function       function:Odorant receptor .,similarity:Belongs to the G-protein coupled in family.,         Background       Olfactory receptor family 1 subfamily L member 8(OR1L8) Homo sapien Olfactory receptors interact with odorant molecules in the nose, to initian neuronal response that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors.	
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms  Observed Band 33kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled in family.,  Background olfactory receptor family 1 subfamily L member 8(OR1L8) Homo sapien Olfactory receptors interact with odorant molecules in the nose, to initian neuronal response that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors (see that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors (see that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors (see that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors (see that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors (see that the protein coupled the protein c	
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 33kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled in family.,  Background olfactory receptor family 1 subfamily L member 8(OR1L8) Homo sapient Olfactory receptors interact with odorant molecules in the nose, to initia neuronal response that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors.	
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band         Observed Band       33kD         Cell Pathway       Cell membrane; Multi-pass membrane protein.         Tissue Specificity       Function         Function       function:Odorant receptor .,similarity:Belongs to the G-protein coupled family.,         Background       olfactory receptor family 1 subfamily L member 8(OR1L8) Homo sapiem Olfactory receptors interact with odorant molecules in the nose, to initial neuronal response that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors (	
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 33kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled in family.,  Background olfactory receptor family 1 subfamily L member 8(OR1L8) Homo sapiem Olfactory receptors interact with odorant molecules in the nose, to initial neuronal response that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors (	
Storage Stability  -20°C/1 year  Synonyms  Observed Band  33kD  Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled a family.,  Background  Olfactory receptor family 1 subfamily L member 8(OR1L8) Homo sapien Olfactory receptors interact with odorant molecules in the nose, to initia neuronal response that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors (	
Synonyms  Observed Band  33kD  Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled in family.,  Background  Olfactory receptor family 1 subfamily L member 8(OR1L8) Homo sapient Olfactory receptors interact with odorant molecules in the nose, to initiat neuronal response that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors (	
Observed Band  Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled in family.,  Olfactory receptor family 1 subfamily L member 8(OR1L8) Homo sapient Olfactory receptors interact with odorant molecules in the nose, to initiat neuronal response that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors (	
Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled a family.,  Background  olfactory receptor family 1 subfamily L member 8(OR1L8) Homo sapient Olfactory receptors interact with odorant molecules in the nose, to initial neuronal response that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors (	
Tissue Specificity  Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled of family.,  Background  olfactory receptor family 1 subfamily L member 8(OR1L8) Homo sapient Olfactory receptors interact with odorant molecules in the nose, to initial neuronal response that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors (	
Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled in family.,  Background  olfactory receptor family 1 subfamily L member 8(OR1L8) Homo sapient Olfactory receptors interact with odorant molecules in the nose, to initial neuronal response that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors (	
family.,  Background  olfactory receptor family 1 subfamily L member 8(OR1L8) Homo sapien Olfactory receptors interact with odorant molecules in the nose, to initia neuronal response that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors (	
Olfactory receptors interact with odorant molecules in the nose, to initial neuronal response that triggers the perception of a smell. The olfactory proteins are members of a large family of G-protein-coupled receptors (	d receptor 1
7-transmembrane domain structure with many neurotransmitter and hor receptors and are responsible for the recognition and G protein-mediate transduction of odorant signals. The olfactory receptor gene family is the genome. The nomenclature assigned to the olfactory receptor gene	tiate a ory receptor os (GPCR) normone ated the largest in

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658








	proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658