



## MGRN1 Polyclonal Antibody

Catalog No	BYab-07752
Isotype	lgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	MGRN1 KIAA0544 RNF156
Protein Name	E3 ubiquitin-protein ligase MGRN1 (EC 6.3.2) (Mahogunin RING finger protein 1) (RING finger protein 156)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	MGRN1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	60kD
Cell Pathway	
Cen Falliway	<ul> <li>Early endosome . The endosomal localization is dependent on the interaction with TSG101.; [Isoform 1]: Cytoplasm, cytosol. Nucleus. Translocation from the cytosol to the nucleus is seen only in the presence of MC1R and MC4R, but not TBXA2R. Excluded from nucleoli.; [Isoform 2]: Cytoplasm, cytosol. Nucleus. Translocation from the cytosol to the nucleus is seen only in the presence of MC1R and MC4R, but not TBXA2R. but not TBXA2R. but not TBXA2R. Excluded from the cytosol to the nucleus is seen only in the presence of MC1R and MC4R, but not TBXA2R. Excluded from nucleoli.; [Isoform 3]: Cytoplasm, cytosol. Cell membrane.; [Isoform 4]: Cytoplasm, cytosol. Cell membrane.</li> </ul>
Tissue Specificity	TSG101.; [Isoform 1]: Cytoplasm, cytosol. Nucleus. Translocation from the cytosol to the nucleus is seen only in the presence of MC1R and MC4R, but not TBXA2R. Excluded from nucleoli.; [Isoform 2]: Cytoplasm, cytosol. Nucleus. Translocation from the cytosol to the nucleus is seen only in the presence of MC1R and MC4R, but not TBXA2R. Excluded from nucleoli.; [Isoform 3]: Cytoplasm, cytosol. Cell membrane.; [Isoform 4]: Cytoplasm, cytosol. Cell
	TSG101.; [Isoform 1]: Cytoplasm, cytosol. Nucleus. Translocation from the cytosol to the nucleus is seen only in the presence of MC1R and MC4R, but not TBXA2R. Excluded from nucleoli.; [Isoform 2]: Cytoplasm, cytosol. Nucleus. Translocation from the cytosol to the nucleus is seen only in the presence of MC1R and MC4R, but not TBXA2R and MC4R, but not TBXA2R. Excluded from nucleoli.; [Isoform 3]: Cytoplasm, cytosol. Cell membrane.; [Isoform 4]: Cytoplasm, cytosol. Cell membrane.

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Background	Mahogunin (MGRN1) is a C3HC4 RING-containing protein with E3 ubiquitin ligase activity in vitro.[supplied by OMIM, Apr 2004],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images** 

Nanjing BYabscience technology Co.,Ltd