



# NMNA2 rabbit pAb

<b>Catalog No</b>	BYab-09848
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	NMNAT2 C1orf15 KIAA0479
<b>Protein Name</b>	Nicotinamide mononucleotide adenylyltransferase 2 (NMN adenylyltransferase 2) (EC 2.7.7.1) (Nicotinate-nucleotide adenylyltransferase 2) (NaMN adenylyltransferase 2) (EC 2.7.7.18)
<b>Immunogen</b>	Synthesized peptide derived from human NMNA2 AA range: 218-268
<b>Specificity</b>	This antibody detects endogenous levels of human NMNA2
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1:1000-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Golgi apparatus membrane ; Lipid-anchor . Cytoplasmic vesicle membrane ; Lipid-anchor . Cytoplasm . Cell projection, axon . Delivered to axons with Golgi-derived cytoplasmic vesicles. .
<b>Tissue Specificity</b>	Highly expressed in brain, in particular in cerebrum, cerebellum, occipital lobe, frontal lobe, temporal lobe and putamen. Also found in the heart, skeletal muscle, pancreas and islets of Langerhans.
<b>Function</b>	
<b>Background</b>	
<b>matters needing attention</b>	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd



**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**