



GFP-Tag Polyclonal Antibody

Catalog No BYab-10281 Isotype IgG Reactivity Species independent Applications WB,IF; Gene Name Formulation Protein Name GFP Tag Immunogen Recombinant Protein of GFP-Tag Specificity GFP antibody is reactive against all variants of Aequorea victoria GFP such as 865T-GFP, RS-GFP, YFP, CFP, RFP and EGFP. Formulation PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using specific immunogen. Dilution WB: 1:5000. IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms YM3206 Observed Band Cell Pathway Tissue Specificity Time green fluorescent protein (GFP) is a protein composed of 238 amino acid residues (26 80/b) that exhibits bright green fluorescence when exposed to light in the blue to ultraviolet range. Although many other marine organisms have similar green fluorescent proteins, GFP traditionally refers to the protein first isolated in peak at a wavelength of 395 nm and a minor one at 475 nm. Its em		
Reactivity Applications WB;IF; Gene Name Protein Name GFP Tag Immunogen Recombinant Protein of GFP-Tag Specificity GFP antibody is reactive against all variants of Aequorea victoria GFP such as S65T-GFP, RS-GFP, YFP, CFP, RFP and EGFP. Formulation PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol. Source Polyclonal, Rabbit.lgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using specific immunogen. Dilution WB: 1:5000. IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms YM3206 Observed Band Cell Pathway Tissue Specificity Function Background The green fluorescent protein (GFP) is a protein composed of 238 amino acid residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultraviolet range. Although many other marine organisms have similar green fluorescent protein first isolated from the jellyfish. The GFP has a major excitation peak at a wavelength of 356 nm and a minor one at 475 nm. Its emission peak at a wavelength of 356 nm and a minor one at 475 nm. Its emission peak at a wavelength of 356 nm and a minor one at 475 nm. Its emission peak is at 500 nm, which is in the lower green	Catalog No	BYab-10281
Applications WB;IF; Gene Name Protein Name GFP Tag Immunogen Recombinant Protein of GFP-Tag Specificity GFP antibody is reactive against all variants of Aequorea victoria GFP such as \$65T-GFP, RS-GFP, YFP, CFP, RFP and EGFP. Formulation PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using specific immunogen. Dilution WB: 1:5000. IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms YM3206 Observed Band Cell Pathway Tissue Specificity Function Background The green fluorescent protein (GFP) is a protein composed of 238 amino acid residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultraviolet range. Although many other marine organisms have similar green fluorescent proteins, GFP traditionally refers to the protein first isolated from the jellyfish. The GFP has a major excitation peak at a wavelength of 395 mm and ammor one at 475 nm. Its emission peak is at 509 nm, which is in the lower green	Isotype	IgG
Gene Name Protein Name GFP Tag Immunogen Recombinant Protein of GFP-Tag Specificity GFP antibody is reactive against all variants of Aequorea victoria GFP such as S65T-GFP, RS-GFP, YFP, CFP, RFP and EGFP. Formulation PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using specific immunogen. Dilution WB: 1:5000. IF 1:50-200 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms YM3206 Observed Band Cell Pathway Tissue Specificity Function Background The green fluorescent protein (GFP) is a protein composed of 238 amino acid residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultraviolet range. Although many other marine organisms have similar green fluorescence proteins, GFP traditionally refers to the protein first isolated from the jellyfish. The GFP has a major excitation peak at a wavelength of 395 nm and aminor one at 475 nm. Its emission peak is at 509 nm, which is in the lower green	Reactivity	Species independent
Protein Name GFP Tag Immunogen Recombinant Protein of GFP-Tag Specificity GFP antibody is reactive against all variants of Aequorea victoria GFP such as S65T-GFP, RS-GFP, YFP, CFP, RFP and EGFP. Formulation PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using specific immunogen. Dilution WB: 1:5000. IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms YM3206 Observed Band Cell Pathway Tissue Specificity Function Background The green fluorescent protein (GFP) is a protein composed of 238 amino acid residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultravolet range. Although many other marine organisms have similar green fluorescent proteins, GFP traditionally refers to the protein first isolated from the jellyfish. The GFP has a major excitation peak at a wavelength of 395 nm and a minor one at 475 nm. Its emission peak is at 509 nm, which is in the lower green	Applications	WB;IF;
Immunogen Recombinant Protein of GFP-Tag Specificity GFP antibody is reactive against all variants of Aequorea victoria GFP such as S65T-GFP, RS-GFP, YFP, CFP, RFP and EGFP. Formulation PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using specific immunogen. Dilution WB: 1:5000. IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms YM3206 Observed Band Cell Pathway Tissue Specificity Function Background The green fluorescent protein (GFP) is a protein composed of 238 amino acid residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultraviolet range. Although many other marine or granisms have similar green fluorescent proteins, GFP traditionally refers to the protein first isolated from the jellyfish. The GFP has a major excitation peak at a wavelength of 395 mm and a minor one at 475 nm. Its emission peak is at 509 mm, which is in the lower green	Gene Name	
Specificity GFP antibody is reactive against all variants of Aequorea victoria GFP such as S65T-GFP, RS-GFP, YFP, CFP, RFP and EGFP. Formulation PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using specific immunogen. Dilution WB: 1:5000. IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms YM3206 Observed Band Cell Pathway Tissue Specificity Function Background The green fluorescent protein (GFP) is a protein composed of 238 amino acid residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultraviolet range. Although many other marine organisms have similar green fluorescent proteins, GFP traditionally refers to the protein first isolated from the jellyfish. The GFP has a major excitation peak at a wavelength of 395 nm and a minor one at 475 nm. Its emission peak is at 509 nm, which is in the lower green	Protein Name	GFP Tag
Formulation PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using specific immunogen. Dilution WB: 1:5000. IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms YM3206 Observed Band Cell Pathway Tissue Specificity Function Background The green fluorescent protein (GFP) is a protein composed of 238 amino acid residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultraviolet range. Although many other marine organisms have similar green fluorescent proteins, GFP traditionally refers to the protein first isolated from the jellyfish. The GFP has a major excitation peak at a wevelength of 395 nm and a minor one at 475 nm. Its emission peak is at 509 nm, which is in the lower green	Immunogen	Recombinant Protein of GFP-Tag
Glycerol. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using specific immunogen. Dilution WB: 1:5000. IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms YM3206 Observed Band Cell Pathway Tissue Specificity Function Background The green fluorescent protein (GFP) is a protein composed of 238 amino acid residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultravolet range. Although many other marine organisms have similar green fluorescent proteins, GFP traditionally refers to the protein first isolated from the jellyfish. The GFP has a major excitation peak at a wavelength of 395 nm and a minor one at 475 nm. Its emission peak is at 509 nm, which is in the lower green	Specificity	GFP antibody is reactive against all variants of Aequorea victoria GFP such as S65T-GFP, RS-GFP, YFP, CFP, RFP and EGFP.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using specific immunogen. Dilution WB: 1:5000. IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms YM3206 Observed Band Cell Pathway Tissue Specificity Function Background The green fluorescent protein (GFP) is a protein composed of 238 amino acid residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultraviolet range. Although many other marine organisms have similar green fluorescent proteins, GFP traditionally refers to the protein first isolated from the jellyfish. The GFP has a major excitation peak at a wavelength of 395 nm and a minor one at 475 nm. Its emission peak is at 509 nm, which is in the lower green	Formulation	
affinity-chromatography using specific immunogen. Dilution WB: 1:5000. IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms YM3206 Observed Band Cell Pathway Tissue Specificity Function Background The green fluorescent protein (GFP) is a protein composed of 238 amino acid residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultraviolet range. Although many other marine organisms have similar green fluorescent proteins, GFP traditionally refers to the protein first isolated from the jellyfish. The GFP has a major excitation peak at a wavelength of 395 nm and a minor one at 475 nm. Its emission peak is at 509 nm, which is in the lower green	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms YM3206 Observed Band Cell Pathway Tissue Specificity Function Background The green fluorescent protein (GFP) is a protein composed of 238 amino acid residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultraviolet range. Although many other marine organisms have similar green fluorescent proteins, GFP traditionally refers to the protein first isolated from the jellyfish. The GFP has a major excitation peak at a wavelength of 395 nm and a minor one at 475 nm. Its emission peak is at 509 nm, which is in the lower green	Purification	·
Purity ≥90% Storage Stability -20°C/1 year Synonyms YM3206 Observed Band Cell Pathway Tissue Specificity Function Background The green fluorescent protein (GFP) is a protein composed of 238 amino acid residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultraviolet range. Although many other marine organisms have similar green fluorescent proteins, GFP traditionally refers to the protein first isolated from the jellyfish. The GFP has a major excitation peak at a wavelength of 395 nm and a minor one at 475 nm. Its emission peak is at 509 nm, which is in the lower green	Dilution	WB: 1:5000. IF 1:50-200
Storage Stability -20°C/1 year Synonyms YM3206 Observed Band Cell Pathway Tissue Specificity Function Background The green fluorescent protein (GFP) is a protein composed of 238 amino acid residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultraviolet range. Although many other marine organisms have similar green fluorescent proteins, GFP traditionally refers to the protein first isolated from the jellyfish. The GFP has a major excitation peak at a wavelength of 395 nm and a minor one at 475 nm. Its emission peak is at 509 nm, which is in the lower green	Concentration	1 mg/ml
Synonyms YM3206 Observed Band Cell Pathway Tissue Specificity Function Background The green fluorescent protein (GFP) is a protein composed of 238 amino acid residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultraviolet range. Although many other marine organisms have similar green fluorescent proteins, GFP has a major excitation peak at a wavelength of 395 nm and a minor one at 475 nm. Its emission peak is at 509 nm, which is in the lower green	Purity	≥90%
Observed Band Cell Pathway Tissue Specificity Function Background The green fluorescent protein (GFP) is a protein composed of 238 amino acid residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultraviolet range. Although many other marine organisms have similar green fluorescent proteins, GFP traditionally refers to the protein first isolated from the jellyfish. The GFP has a major excitation peak at a wavelength of 395 nm and a minor one at 475 nm. Its emission peak is at 509 nm, which is in the lower green	Storage Stability	-20°C/1 year
Cell Pathway Tissue Specificity Function Background The green fluorescent protein (GFP) is a protein composed of 238 amino acid residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultraviolet range. Although many other marine organisms have similar green fluorescent proteins, GFP traditionally refers to the protein first isolated from the jellyfish. The GFP has a major excitation peak at a wavelength of 395 nm and a minor one at 475 nm. Its emission peak is at 509 nm, which is in the lower green	Synonyms	YM3206
Tissue Specificity Function Background The green fluorescent protein (GFP) is a protein composed of 238 amino acid residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultraviolet range. Although many other marine organisms have similar green fluorescent proteins, GFP traditionally refers to the protein first isolated from the jellyfish. The GFP has a major excitation peak at a wavelength of 395 nm and a minor one at 475 nm. Its emission peak is at 509 nm, which is in the lower green	Observed Band	
Function Background The green fluorescent protein (GFP) is a protein composed of 238 amino acid residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultraviolet range. Although many other marine organisms have similar green fluorescent proteins, GFP traditionally refers to the protein first isolated from the jellyfish. The GFP has a major excitation peak at a wavelength of 395 nm and a minor one at 475 nm. Its emission peak is at 509 nm, which is in the lower green	Cell Pathway	
Background The green fluorescent protein (GFP) is a protein composed of 238 amino acid residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultraviolet range. Although many other marine organisms have similar green fluorescent proteins, GFP traditionally refers to the protein first isolated from the jellyfish. The GFP has a major excitation peak at a wavelength of 395 nm and a minor one at 475 nm. Its emission peak is at 509 nm, which is in the lower green	Tissue Specificity	
residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultraviolet range. Although many other marine organisms have similar green fluorescent proteins, GFP traditionally refers to the protein first isolated from the jellyfish. The GFP has a major excitation peak at a wavelength of 395 nm and a minor one at 475 nm. Its emission peak is at 509 nm, which is in the lower green	Function	
	Background	residues (26.9kD) that exhibits bright green fluorescence when exposed to light in the blue to ultraviolet range. Although many other marine organisms have similar green fluorescent proteins, GFP traditionally refers to the protein first isolated from the jellyfish. The GFP has a major excitation peak at a wavelength of 395 nm and a minor one at 475 nm. Its emission peak is at 509 nm, which is in the lower green

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658






matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



1ug GFP fusion protein+ Primary antibody dilution at 1) 1:5000 2) 1:20000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658