



TRAF1 Polyclonal Antibody

Catalog No	BYab-10596
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	TRAF1
Protein Name	TRAF1
Immunogen	Synthesized peptide derived from TRAF1 at AA range: 191-240
Specificity	TRAF1 Polyclonal Antibody detects endogenous levels of TRAF1
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000, ELISA 1:10000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	TNF receptor-associated factor 1 (Epstein-Barr virus-induced protein 6)
Observed Band	46kD
Cell Pathway	cytoplasm,cytosol,
Tissue Specificity	Lymphoma,Skin,Spleen,Stomach,
Function	domain:The coiled coil domain mediates homo- and hetero-oligomerization.,domain:The MATH/TRAF domain binds to receptor cytoplasmic domains.,function:Adapter protein and signal transducer that links members of the tumor necrosis factor receptor family to different signaling pathways by association with the receptor cytoplasmic domain and kinases. Mediates activation of NF-kappa-B and JNK and is involved in apoptosis. The TRAF1/TRAF2 complex recruits the apoptotic suppressors BIRC2 and BIRC3 to TNFRSF1B/TNFR2.,similarity:Contains 1 MATH domain.,subunit:Homotrimer (Probable). Heteromer with TRAF2 and associates with TNFRSF1B/TNFR2 through TRAF2. Associates with TNFRSF4, TNFRSF5/CD40, TNFRSF8/CD30, TNFRSF9/CD137, TNFRSF11A/RANK, TNFRSF18/AITR, TNFRSF17/BCMA, TNFRSF19/TROY, TNFRSF19L/RELT, XEDAR, EDAR, Epstein-Barr virus

Nanjing BYabscience technology Co.,Ltd



BNFL1/LMP-1, TANK/ITRAF, TRAF1 and RIPK2. Interacts with BIRC2 and BIRC3

Background

TNF receptor associated factor 1 (TRAF1) Homo sapiens The protein encoded by this gene is a member of the TNF receptor (TNFR) associated factor (TRAF) protein family. TRAF proteins associate with, and mediate the signal transduction from various receptors of the TNFR superfamily. This protein and TRAF2 form a heterodimeric complex, which is required for TNF-alpha-mediated activation of MAPK8/JNK and NF-kappaB. The protein complex formed by this protein and TRAF2 also interacts with inhibitor-of-apoptosis proteins (IAPs), and thus mediates the anti-apoptotic signals from TNF receptors. The expression of this protein can be induced by Epstein-Barr virus (EBV). EBV infection membrane protein 1 (LMP1) is found to interact with this and other TRAF proteins; this interaction is thought to link LMP1-mediated B lymphocyte transformation to the signal transduction from TNFR family receptors. Three transcript variants encoding two different isoforms have


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of HEPG2 cells using TRAF1 Polyclonal Antibody diluted at 1:500. Secondary antibody(catalog#:RS0002) was diluted at 1:20000

Nanjing BYabs science technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658