



Prolactin Polyclonal Antibody

Catalog No BYab-16047 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB;IHC;IF;ELISA Gene Name PRL Protein Name Prolactin Immunogen The antiserum was produced against synthesized peptide derived from the Internal region of human PRL. AA range:131-180 Specificity Prolactin Polyclonal Antibody detects endogenous levels of Prolactin protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/20000. IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms PRL; Prolactin; PRL Observed Band 25kD Cell Pathway Secreted. Tissue Specificity Mammary gland, Pituitary, Testis, Function function:Prolactin acts primarily on the mammary gland by promoting lacitation, online information:Prolactin entry, similarity: Bel		
Reactivity Human;Rat;Mouse; Applications WB;IHC;IF;ELISA Gene Name PRL Protein Name Prolactin Immunogen The antiserum was produced against synthesized peptide derived from the Internal region of human PRL. AA range:131-180 Specificity Prolactin Polyclonal Antibody detects endogenous levels of Prolactin protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms PRL; Prolactin; PRL Observed Band 25kD Cell Pathway Secreted. Tissue Specificity Mammary gland,Pituitary,Testis, Function function:Prolactin acts primarily on the mammary gland by promoting lactation, online information: Prolactin entry, similarity; Belongs to the somatotropin/prolactin family., Background This gene encodes the anterior pituitary hormone prolactin. This secreted hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing apoptosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants	Catalog No	BYab-16047
Applications WB;IHC;IF;ELISA Gene Name PRL Protein Name Prolactin Immunogen The antiserum was produced against synthesized peptide derived from the Internal region of human PRL. AA range:131-180 Specificity Prolactin Polyclonal Antibody detects endogenous levels of Prolactin protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/20000. IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms PRL; Prolactin; PRL Observed Band 25kD Cell Pathway Secreted. Tissue Specificity Mammary gland, Pituitary, Testis, Function function: Prolactin acts primarily on the mammary gland by promoting lactation, online information: Prolactin entry, similarity: Belongs to the somatotropin/prolactin family., Background This gene encodes the anterior pituitary hormone prolactin. This secreted hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing appoptosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants	Isotype	IgG
Gene Name PRL Protein Name Prolactin Immunogen The antiserum was produced against synthesized peptide derived from the Internal region of human PRL. AA range:131-180 Specificity Prolactin Polyclonal Antibody detects endogenous levels of Prolactin protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms PRL; Prolactin; PRL Observed Band 25kD Cell Pathway Secreted. Tissue Specificity Mammary gland, Pituitary, Testis, Function function: Prolactin acts primarily on the mammary gland by promoting lactation, online information: Prolactin entry, similarity: Belongs to the somatotropin/prolactin family., Background This gene encodes the anterior pituitary hormone prolactin. This secreted hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing apoptosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants	Reactivity	Human;Rat;Mouse;
Protein Name Prolactin Immunogen The antiserum was produced against synthesized peptide derived from the Internal region of human PRL. AA range:131-180 Specificity Prolactin Polyclonal Antibody detects endogenous levels of Prolactin protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms PRL; Prolactin; PRL Observed Band 25KD Cell Pathway Secreted. Tissue Specificity Mammary gland, Pituitary, Testis, Function function: Prolactin acts primarily on the mammary gland by promoting lactation, online information: Prolactin entry, similarity: Belongs to the somatotropin/prolactin family. Background This gene encodes the anterior pituitary hormone prolactin. This secreted hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing apoptosis, and it is essential for lactation.	Applications	WB;IHC;IF;ELISA
Immunogen The antiserum was produced against synthesized peptide derived from the Internal region of human PRL. AA range:131-180 Specificity Prolactin Polyclonal Antibody detects endogenous levels of Prolactin protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/20000. IF 1:50-200 Concentration 1 mg/ml Purity 990% Storage Stability -20°C/1 year Synonyms PRL; Prolactin; PRL Observed Band 25kD Cell Pathway Secreted. Tissue Specificity Mammary gland, Pituitary, Testis, Function function: Prolactin acts primarily on the mammary gland by promoting lactation., online information: Prolactin entry, similarity: Belongs to the somatotropin/prolactin family., Background This gene encodes the anterior pituitary hormone prolactin. This secreted hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing apoptosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants	Gene Name	PRL
Internal region of human PRL. AA range:131-180 Specificity Prolactin Polyclonal Antibody detects endogenous levels of Prolactin protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms PRL; Prolactin; PRL Observed Band 25kD Cell Pathway Secreted. Tissue Specificity Mammary gland, Pituitary, Testis, Function function: Prolactin acts primarily on the mammary gland by promoting lactation, online information: Prolactin entry, similarity: Belongs to the somatotropin/prolactin family., Background This gene encodes the anterior pituitary hormone prolactin. This secreted hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing apoptosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants	Protein Name	Prolactin
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms PRL; Prolactin; PRL Observed Band 25kD Cell Pathway Secreted. Tissue Specificity Mammary gland, Pituitary, Testis, Function function: Prolactin acts primarily on the mammary gland by promoting lactation., online information: Prolactin entry, similarity: Belongs to the somatotropin/prolactin family. Background This gene encodes the anterior pituitary hormone prolactin. This secreted hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing appotosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants	Immunogen	
Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms PRL; Prolactin; PRL Observed Band 25kD Cell Pathway Secreted. Tissue Specificity Mammary gland, Pituitary, Testis, Function function: Prolactin acts primarily on the mammary gland by promoting lactation., online information: Prolactin entry, similarity: Belongs to the somatotropin/prolactin family Background This gene encodes the anterior pituitary hormone prolactin. This secreted hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing apoptosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants	Specificity	Prolactin Polyclonal Antibody detects endogenous levels of Prolactin protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms PRL; Prolactin; PRL Observed Band 25kD Cell Pathway Secreted. Tissue Specificity Mammary gland, Pituitary, Testis, Function function: Prolactin acts primarily on the mammary gland by promoting lactation., online information: Prolactin entry, similarity: Belongs to the somatotropin/prolactin family., Background This gene encodes the anterior pituitary hormone prolactin. This secreted hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing apoptosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/20000 IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms PRL; Prolactin; PRL Observed Band 25kD Cell Pathway Secreted. Tissue Specificity Mammary gland, Pituitary, Testis, Function function: Prolactin acts primarily on the mammary gland by promoting lactation., online information: Prolactin entry, similarity: Belongs to the somatotropin/prolactin family., Background This gene encodes the anterior pituitary hormone prolactin. This secreted hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing apoptosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms PRL; Prolactin; PRL Observed Band 25kD Cell Pathway Secreted. Tissue Specificity Mammary gland,Pituitary,Testis, Function function:Prolactin acts primarily on the mammary gland by promoting lactation.,online information:Prolactin entry,similarity:Belongs to the somatotropin/prolactin family., Background This gene encodes the anterior pituitary hormone prolactin. This secreted hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing apoptosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms PRL; Prolactin; PRL Observed Band 25kD Cell Pathway Secreted. Tissue Specificity Mammary gland,Pituitary,Testis, Function function:Prolactin acts primarily on the mammary gland by promoting lactation.,online information:Prolactin entry,similarity:Belongs to the somatotropin/prolactin family., Background This gene encodes the anterior pituitary hormone prolactin. This secreted hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing apoptosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants	Dilution	WB: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/20000 IF 1:50-200
Storage Stability -20°C/1 year PRL; Prolactin; PRL Observed Band 25kD Cell Pathway Secreted. Tissue Specificity Mammary gland, Pituitary, Testis, Function function: Prolactin acts primarily on the mammary gland by promoting lactation., online information: Prolactin entry, similarity: Belongs to the somatotropin/prolactin family., Background This gene encodes the anterior pituitary hormone prolactin. This secreted hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing apoptosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants	Concentration	1 mg/ml
Synonyms PRL; Prolactin; PRL Observed Band 25kD Cell Pathway Secreted. Tissue Specificity Mammary gland, Pituitary, Testis, Function function: Prolactin acts primarily on the mammary gland by promoting lactation., online information: Prolactin entry, similarity: Belongs to the somatotropin/prolactin family., Background This gene encodes the anterior pituitary hormone prolactin. This secreted hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing apoptosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants	Purity	≥90%
Observed Band Cell Pathway Secreted. Tissue Specificity Mammary gland, Pituitary, Testis, Function function: Prolactin acts primarily on the mammary gland by promoting lactation., online information: Prolactin entry, similarity: Belongs to the somatotropin/prolactin family., Background This gene encodes the anterior pituitary hormone prolactin. This secreted hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing apoptosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants	Storage Stability	-20°C/1 year
Cell Pathway Secreted. Tissue Specificity Mammary gland, Pituitary, Testis, Function function: Prolactin acts primarily on the mammary gland by promoting lactation., online information: Prolactin entry, similarity: Belongs to the somatotropin/prolactin family., Background This gene encodes the anterior pituitary hormone prolactin. This secreted hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing apoptosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants	Synonyms	PRL; Prolactin; PRL
Tissue Specificity Mammary gland,Pituitary,Testis, function function:Prolactin acts primarily on the mammary gland by promoting lactation.,online information:Prolactin entry,similarity:Belongs to the somatotropin/prolactin family., Background This gene encodes the anterior pituitary hormone prolactin. This secreted hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing apoptosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants	Observed Band	25kD
Function function:Prolactin acts primarily on the mammary gland by promoting lactation.,online information:Prolactin entry,similarity:Belongs to the somatotropin/prolactin family., Background This gene encodes the anterior pituitary hormone prolactin. This secreted hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing apoptosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants	Cell Pathway	Secreted.
lactation.,online information:Prolactin entry,similarity:Belongs to the somatotropin/prolactin family., Background This gene encodes the anterior pituitary hormone prolactin. This secreted hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing apoptosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants	Tissue Specificity	Mammary gland,Pituitary,Testis,
hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing apoptosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants	Function	lactation.,online information:Prolactin entry,similarity:Belongs to the
	Background	hormone is a growth regulator for many tissues, including cells of the immune system. It may also play a role in cell survival by suppressing apoptosis, and it is essential for lactation. Alternative splicing results in multiple transcript variants

Nanjing BYabscience technology Co.,Ltd






matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of SKOV3 cells using Prolactin Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:100

Nanjing BYabscience technology Co.,Ltd