




## GBP3 Polyclonal Antibody

Catalog No         BYab-16160           Isotype         IgG           Reactivity         Human;Rat;Mouse;           Applications         WB;IHC;IF;ELISA           Gene Name         GBP3           Protein Name         Guanylate-binding protein 3           Immunogen         The antiserum was produced against synthesized peptide derived from human GBP3, AA range:481-530           Specificity         GBP3 Polyclonal Antibody detects endogenous levels of GBP3 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Polyclonal, Rabbit,IgG           Purification         The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/10000 IF 1:50-200           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         GBP3; Guanylate-binding protein 3; GTP-binding protein 3; GBP-3; Guanine nucleotide-binding protein 3           Observed Band         64kD           Cell Pathway         Cytoplasm. Cytoplasm. perinuclear region. Golgi apparatus membrane Heterodimers with GBP1; GBP2 and GBP5 localize in the compartment of the protein swith GBP1; and the Golgi apparatus membrane Potential poly-A sequence. sequence caution:Unl		
Reactivity Human;Rat;Mouse;  Applications WB;IHC;IF;ELISA  Gene Name GBP3  Protein Name Guanylate-binding protein 3  Immunogen The antiserum was produced against synthesized peptide derived from human GBP3. AA range:481-530  Specificity GBP3 Polyclonal Antibody detects endogenous levels of GBP3 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/10000 IF 1:50-200  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms GBP3; Guanylate-binding protein 3; GTP-binding protein 3; GBP-3; Guanine nucleotide-binding protein 3  Observed Band 64kD  Cell Pathway Cytoplasm. Cytoplasm, perinuclear region. Golgi apparatus membrane. Heterodimers with GBP1, GBP2 and GBP5 localize in the compartment of the prenylated GBPs: with GBP1 in a vesicle-like compartment, with GBP2, around the nucleus and with GBP-5, at the Golgi apparatus.  Tissue Specificity Brain,Prostate,  Function function:Binds GTP, GDP and GMP, sequence caution:Contaminating sequence. Potential poly-A sequence, sequence caution:Unlikely isoform. Aberrant splice sites.,similarity:Belongs to the GBP family.  GBPs specifically bind guanine nucleotides (GMP, GDP, and GTP) and COntain to Yof the three consensus motifs found in typical GTP-binding protein. The	Catalog No	BYab-16160
Applications WB;IHC;IF;ELISA  Gene Name GBP3  Protein Name Guanylate-binding protein 3  Immunogen The antiserum was produced against synthesized peptide derived from human GBP3. AA range:481-530  Specificity GBP3 Polyclonal Antibody detects endogenous levels of GBP3 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/10000 IF 1:50-200  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms GBP3; Guanylate-binding protein 3; GTP-binding protein 3; GBP-3; Guanine nucleotide-binding protein 3  Observed Band 64kD  Cell Pathway Cytoplasm. Cytoplasm. perinuclear region. Golgi apparatus membrane. Heterodimers with GBP1, GBP2 and GBP5 localize in the compartment of the prenylated GBPs: with GBP1 in a vesicle-like compartment, with GBP2, around the nucleus and with GBP-5, at the Golgi apparatus.  Tissue Specificity Brain, Prostate,  Function function: Binds GTP, GDP and GMP, sequence caution: Contaminating sequence. Potential poly-A sequence. sequence caution: Unlikely isoform. Aberrant splice sites., similarity: Belongs to the GBP family., GBPs specifically bind guanine nucleotides (GMP, GDP, and GTP), and GDP, and GD	Isotype	IgG
Protein Name GBP3 Protein Name Guanylate-binding protein 3  Immunogen The antiserum was produced against synthesized peptide derived from human GBP3. AA range:481-530  Specificity GBP3 Polyclonal Antibody detects endogenous levels of GBP3 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/10000 IF 1:50-200  Concentration 1 mg/ml  Purity 290%  Storage Stability -20°C/1 year  Synonyms GBP3; Guanylate-binding protein 3; GTP-binding protein 3; GBP-3; Guanine nucleotide-binding protein 3  Observed Band 64kD  Cell Pathway Cytoplasm. Cytoplasm, perinuclear region. Golgi apparatus membrane. Heterodimers with GBP1, GBP2 and GBP5 localize in the compartment of the prenylated GBPs: with GBP-1 in a vesicle-like compartment, with GBP2, around the nucleus and with GBP-5, at the Golgi apparatus.  Tissue Specificity Brain, Prostate,  Function function: Binds GTP, GDP and GMP., sequence caution: Contaminating sequence. Potential poly-A sequence, sequence caution: Unlikely isoform. Aberrant splice sites, similanty. Belongs to the GBP family.  GBPs specifically bind guanine nucleotides (GMP, GDP, and GTP) and contain two of the three consensus motifs found in typical GTP-binding proteins. The	Reactivity	Human;Rat;Mouse;
Protein Name         Guanylate-binding protein 3           Immunogen         The antiserum was produced against synthesized peptide derived from human GBP3. AA range:481-530           Specificity         GBP3 Polyclonal Antibody detects endogenous levels of GBP3 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Polyclonal, Rabbit,IgG           Purification         The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/10000 IF 1:50-200           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         GBP3; Guanylate-binding protein 3; GTP-binding protein 3; GBP-3; Guanine nucleotide-binding protein 3           Observed Band         64kD           Cell Pathway         Cytoplasm . Cytoplasm , perinuclear region . Golgi apparatus membrane . Heterodimers with GBP1 / GBP2 and GBP5 localize in the compartment of the prenylated GBPs: with GBP1 in a vesicle-like compartment, with GBP2, around the nucleus and with GBP-5, at the Golgi apparatus           Tissue Specificity         Brain,Prostate,           Function         function:Binds GTP, GDP and GMP., sequence caution:Contaminating sequence. Potential poly-A sequence, sequence caution:Unlikely isoform. Aberrant splice sites, similarity:Belongs to th	Applications	WB;IHC;IF;ELISA
Immunogen  The antiserum was produced against synthesized peptide derived from human GBP3. AA range:481-530  Specificity  GBP3 Polyclonal Antibody detects endogenous levels of GBP3 protein.  Formulation  Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source  Polyclonal, Rabbit,IgG  Purification  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/10000 IF 1:50-200  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  GBP3; Guanylate-binding protein 3; GTP-binding protein 3; GBP-3; Guanine nucleotide-binding protein 3  Observed Band  64kD  Cell Pathway  Cytoplasm . Cytoplasm, perinuclear region . Golgi apparatus membrane . Heterodimers with GBP1, GBP2 and GBP5 localize in the compartment of the prenylated GBPs: with GBP1 in a vesicle-like compartment, with GBP2, around the nucleus and with GBP-5, at the Golgi apparatus.  Tissue Specificity  Brain,Prostate,  Function  function:Binds GTP, GDP and GMP., sequence caution:Contaminating sequence. Potential poly-A sequence, sequence caution:Unlikely isoform. Aberrant splice sites, similarity:Belongs to the GBP family.  GBPs specifically bind guanine nucleotides (GMP, GDP, and GTP) and contain two of the three consensus motifs found in typical GTP) and contain two of the three consensus motifs found in typical GTP) and contain two of the three consensus motifs found in typical GTP) and contain two of the three consensus motifs found in typical GTP) and contain two of the three consensus motifs found in typical GTP) and contain two of the three consensus motifs found in typical GTP) and contain two of the three consensus motifs found in typical GTP) and contain two of the three consensus motifs found in typical GTP-binding proteins. The	Gene Name	GBP3
Specificity GBP3 Polyclonal Antibody detects endogenous levels of GBP3 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/10000 IF 1:50-200  Concentration 1 mg/ml  Purity 290%  Storage Stability -20°C/1 year  Synonyms GBP3; Guanylate-binding protein 3; GTP-binding protein 3; GBP-3; Guanine nucleotide-binding protein 3  Observed Band 64kD  Cell Pathway Cytoplasm. Cytoplasm, perinuclear region. Golgi apparatus membrane. Heterodimers with GBP1, GBP2 and GBP5 localize in the compartment of the prenylated GBPs: with GBP1 in a vesicle-like compartment, with GBP2, around the nucleus and with GBP-5, at the Golgi apparatus.  Tissue Specificity Brain, Prostate,  Function function: Binds GTP, GDP and GMP, sequence caution: Contaminating sequence. Potential poly-A sequence, sequence caution: Unlikely isoform. Aberrant splice sites, similarity: Belongs to the GBP family.  Background This gene encodes a member of the guanylate-binding protein (GBP) family. GBPs specifically bind guannine nucleotides (GMP, GDP, and GTP-binding proteins. The	Protein Name	Guanylate-binding protein 3
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/10000 IF 1:50-200  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms GBP3; Guanylate-binding protein 3; GTP-binding protein 3; GBP-3; Guanine nucleotide-binding protein 3  Observed Band 64kD  Cytoplasm . Cytoplasm, perinuclear region . Golgi apparatus membrane . Heterodimers with GBP1, GBP2 and GBP5 localize in the compartment of the prenylated GBPs: with GBP1 in a vesicle-like compartment, with GBP2, around the nucleus and with GBP-5, at the Golgi apparatus  Tissue Specificity Brain, Prostate,  Function function:Binds GTP, GDP and GMPsequence caution:Contaminating sequence. Potential poly-A sequencesequence caution:Unlikely isoform. Aberrant splice sitessimilarity:Belongs to the GBP family.,  Background This gene encodes a member of the guanylate-binding protein (GBP) family. GBPs specifically bind quannine nucleotides (GMP, GDP, and GTP)-and GTP-binding proteins. The	Immunogen	
Source         Polyclonal, Rabbit, IgG           Purification         The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/10000 IF 1:50-200           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         GBP3; Guanylate-binding protein 3; GTP-binding protein 3; GBP-3; Guanine nucleotide-binding protein 3           Observed Band         64kD           Cell Pathway         Cytoplasm . Cytoplasm, perinuclear region . Golgi apparatus membrane . Heterodimers with GBP1, GBP2 and GBP5 localize in the compartment of the prenylated GBPs: with GBP1 in a vesicle-like compartment, with GBP2, around the nucleus and with GBP-5, at the Golgi apparatus .           Tissue Specificity         Brain,Prostate,           Function         function:Binds GTP, GDP and GMP, sequence caution:Contaminating sequence. Potential poly-A sequence, sequence caution:Unlikely isoform. Aberrant splice sites., similarity:Belongs to the GBP family.           Background         This gene encodes a member of the guanylate-binding protein (GBP) family. GBPs specifically bind guanine nucleotides (GMP, GDP, and GTP) and contain two of the three consensus motifs found in typical GTP-binding proteins. The	Specificity	GBP3 Polyclonal Antibody detects endogenous levels of GBP3 protein.
Purification  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/10000 IF 1:50-200  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  GBP3; Guanylate-binding protein 3; GTP-binding protein 3; GBP-3; Guanine nucleotide-binding protein 3  Observed Band  64kD  Cell Pathway  Cytoplasm . Cytoplasm, perinuclear region . Golgi apparatus membrane . Heterodimers with GBP1, GBP2 and GBP5 localize in the compartment of the prenylated GBPs: with GBP1 in a vesicle-like compartment, with GBP2, around the nucleus and with GBP-5, at the Golgi apparatus.  Tissue Specificity  Brain,Prostate,  Function  function:Binds GTP, GDP and GMP., sequence caution:Contaminating sequence. Potential poly-A sequence., sequence caution:Unlikely isoform. Aberrant splice sites., similarity:Belongs to the GBP family.  Background  This gene encodes a member of the guanylate-binding protein (GBP) family. GBPs specifically bind guanine nucleotides (GMP, GDP, and GTP) and contain two of the three consensus motifs found in typical GTP-binding proteins. The	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/10000 IF 1:50-200  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms GBP3; Guanylate-binding protein 3; GTP-binding protein 3; GBP-3; Guanine nucleotide-binding protein 3  Observed Band 64kD  Cell Pathway Cytoplasm . Cytoplasm, perinuclear region . Golgi apparatus membrane . Heterodimers with GBP1, GBP2 and GBP5 localize in the compartment of the prenylated GBPs: with GBP1 in a vesicle-like compartment, with GBP2, around the nucleus and with GBP-5, at the Golgi apparatus  Tissue Specificity Brain,Prostate,  Function function:Binds GTP, GDP and GMP.,sequence caution:Contaminating sequence. Potential poly-A sequence.,sequence caution:Unlikely isoform. Aberrant splice sites.,similarity:Belongs to the GBP family.,  Background This gene encodes a member of the guanylate-binding protein (GBP) family. GBPs specifically bind guanine nucleotides (GMP, GDP, and GTP) and contain two of the three consensus motifs found in typical GTP-binding proteins. The	Source	Polyclonal, Rabbit,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       GBP3; Guanylate-binding protein 3; GTP-binding protein 3; GBP-3; Guanine nucleotide-binding protein 3         Observed Band       64kD         Cell Pathway       Cytoplasm . Cytoplasm, perinuclear region . Golgi apparatus membrane . Heterodimers with GBP1, GBP2 and GBP5 localize in the compartment of the prenylated GBPs: with GBP1 in a vesicle-like compartment, with GBP2, around the nucleus and with GBP-5, at the Golgi apparatus.         Tissue Specificity       Brain,Prostate,         Function       function:Binds GTP, GDP and GMP.,sequence caution:Contaminating sequence. Potential poly-A sequence.,sequence caution:Unlikely isoform. Aberrant splice sites, similarity:Belongs to the GBP family.         Background       This gene encodes a member of the guanylate-binding protein (GBP) family. GBPs specifically bind guanine nucleotides (GMP, GDP, and GTP) and contain two of the three consensus motifs found in typical GTP-binding proteins. The	Purification	·
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms GBP3; Guanylate-binding protein 3; GTP-binding protein 3; GBP-3; Guanine nucleotide-binding protein 3  Observed Band 64kD  Cell Pathway Cytoplasm . Cytoplasm , perinuclear region . Golgi apparatus membrane . Heterodimers with GBP1, GBP2 and GBP5 localize in the compartment of the prenylated GBPs: with GBP1 in a vesicle-like compartment, with GBP2, around the nucleus and with GBP-5, at the Golgi apparatus  Tissue Specificity Brain,Prostate,  Function function:Binds GTP, GDP and GMP.,sequence caution:Contaminating sequence. Potential poly-A sequence.,sequence caution:Unlikely isoform. Aberrant splice sites.,similarity:Belongs to the GBP family.,  Background This gene encodes a member of the guanylate-binding protein (GBP) family. GBPs specifically bind guanine nucleotides (GMP, GDP, and GTP) and contain two of the three consensus motifs found in typical GTP-binding proteins. The	Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/10000 IF 1:50-200
Storage Stability  -20°C/1 year  GBP3; Guanylate-binding protein 3; GTP-binding protein 3; GBP-3; Guanine nucleotide-binding protein 3  Observed Band  64kD  Cell Pathway  Cytoplasm . Cytoplasm, perinuclear region . Golgi apparatus membrane . Heterodimers with GBP1, GBP2 and GBP5 localize in the compartment of the prenylated GBPs: with GBP1 in a vesicle-like compartment, with GBP2, around the nucleus and with GBP-5, at the Golgi apparatus  Tissue Specificity  Brain,Prostate,  Function  function:Binds GTP, GDP and GMP., sequence caution:Contaminating sequence. Potential poly-A sequence., sequence caution:Unlikely isoform. Aberrant splice sites., similarity:Belongs to the GBP family.,  This gene encodes a member of the guanylate-binding protein (GBP) family. GBPs specifically bind guanine nucleotides (GMP, GDP, and GTP) and contain two of the three consensus motifs found in typical GTP-binding proteins. The		1 ma/ml
Synonyms  GBP3; Guanylate-binding protein 3; GTP-binding protein 3; GBP-3; Guanine nucleotide-binding protein 3  Observed Band  64kD  Cell Pathway  Cytoplasm . Cytoplasm, perinuclear region . Golgi apparatus membrane . Heterodimers with GBP1, GBP2 and GBP5 localize in the compartment of the prenylated GBPs: with GBP1 in a vesicle-like compartment, with GBP2, around the nucleus and with GBP-5, at the Golgi apparatus  Tissue Specificity  Brain,Prostate,  Function  function:Binds GTP, GDP and GMP.,sequence caution:Contaminating sequence. Potential poly-A sequence.,sequence caution:Unlikely isoform. Aberrant splice sites.,similarity:Belongs to the GBP family.,  This gene encodes a member of the guanylate-binding protein (GBP) family. GBPs specifically bind guanine nucleotides (GMP, GDP, and GTP) and contain two of the three consensus motifs found in typical GTP-binding proteins. The	Concentration	i ilig/ilii
nucleotide-binding protein 3  Observed Band  Cytoplasm . Cytoplasm, perinuclear region . Golgi apparatus membrane . Heterodimers with GBP1, GBP2 and GBP5 localize in the compartment of the prenylated GBPs: with GBP1 in a vesicle-like compartment, with GBP2, around the nucleus and with GBP-5, at the Golgi apparatus  Tissue Specificity  Brain,Prostate,  Function  function:Binds GTP, GDP and GMP.,sequence caution:Contaminating sequence. Potential poly-A sequence.,sequence caution:Unlikely isoform. Aberrant splice sites.,similarity:Belongs to the GBP family.,  This gene encodes a member of the guanylate-binding protein (GBP) family. GBPs specifically bind guanine nucleotides (GMP, GDP, and GTP) and contain two of the three consensus motifs found in typical GTP-binding proteins. The		
Cytoplasm . Cytoplasm, perinuclear region . Golgi apparatus membrane . Heterodimers with GBP1, GBP2 and GBP5 localize in the compartment of the prenylated GBPs: with GBP1 in a vesicle-like compartment, with GBP2, around the nucleus and with GBP-5, at the Golgi apparatus  Tissue Specificity  Brain,Prostate,  Function  function:Binds GTP, GDP and GMP.,sequence caution:Contaminating sequence. Potential poly-A sequence.,sequence caution:Unlikely isoform. Aberrant splice sites.,similarity:Belongs to the GBP family.,  This gene encodes a member of the guanylate-binding protein (GBP) family. GBPs specifically bind guanine nucleotides (GMP, GDP, and GTP) and contain two of the three consensus motifs found in typical GTP-binding proteins. The	Purity	≥90%
Heterodimers with GBP1, GBP2 and GBP5 localize in the compartment of the prenylated GBPs: with GBP1 in a vesicle-like compartment, with GBP2, around the nucleus and with GBP-5, at the Golgi apparatus.  Tissue Specificity  Brain,Prostate,  Function  function:Binds GTP, GDP and GMP.,sequence caution:Contaminating sequence. Potential poly-A sequence.,sequence caution:Unlikely isoform. Aberrant splice sites.,similarity:Belongs to the GBP family.,  This gene encodes a member of the guanylate-binding protein (GBP) family. GBPs specifically bind guanine nucleotides (GMP, GDP, and GTP) and contain two of the three consensus motifs found in typical GTP-binding proteins. The	Purity Storage Stability	≥90% -20°C/1 year GBP3; Guanylate-binding protein 3; GTP-binding protein 3; GBP-3; Guanine
Function  function:Binds GTP, GDP and GMP.,sequence caution:Contaminating sequence. Potential poly-A sequence.,sequence caution:Unlikely isoform. Aberrant splice sites.,similarity:Belongs to the GBP family.,  This gene encodes a member of the guanylate-binding protein (GBP) family. GBPs specifically bind guanine nucleotides (GMP, GDP, and GTP) and contain two of the three consensus motifs found in typical GTP-binding proteins. The	Purity Storage Stability Synonyms	≥90% -20°C/1 year  GBP3; Guanylate-binding protein 3; GTP-binding protein 3; GBP-3; Guanine nucleotide-binding protein 3
Potential poly-A sequence., sequence caution: Unlikely isoform. Aberrant splice sites., similarity: Belongs to the GBP family.,  Background  This gene encodes a member of the guanylate-binding protein (GBP) family.  GBPs specifically bind guanine nucleotides (GMP, GDP, and GTP) and contain two of the three consensus motifs found in typical GTP-binding proteins. The	Purity Storage Stability Synonyms Observed Band	≥90%  -20°C/1 year  GBP3; Guanylate-binding protein 3; GTP-binding protein 3; GBP-3; Guanine nucleotide-binding protein 3  64kD  Cytoplasm . Cytoplasm, perinuclear region . Golgi apparatus membrane . Heterodimers with GBP1, GBP2 and GBP5 localize in the compartment of the prenylated GBPs: with GBP1 in a vesicle-like compartment, with GBP2, around
GBPs specifically bind guanine nucleotides (GMP, GDP, and GTP) and contain two of the three consensus motifs found in typical GTP-binding proteins. The	Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90%  -20°C/1 year  GBP3; Guanylate-binding protein 3; GTP-binding protein 3; GBP-3; Guanine nucleotide-binding protein 3  64kD  Cytoplasm . Cytoplasm, perinuclear region . Golgi apparatus membrane . Heterodimers with GBP1, GBP2 and GBP5 localize in the compartment of the prenylated GBPs: with GBP1 in a vesicle-like compartment, with GBP2, around the nucleus and with GBP-5, at the Golgi apparatus
	Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90%  -20°C/1 year  GBP3; Guanylate-binding protein 3; GTP-binding protein 3; GBP-3; Guanine nucleotide-binding protein 3  64kD  Cytoplasm . Cytoplasm, perinuclear region . Golgi apparatus membrane . Heterodimers with GBP1, GBP2 and GBP5 localize in the compartment of the prenylated GBPs: with GBP1 in a vesicle-like compartment, with GBP2, around the nucleus and with GBP-5, at the Golgi apparatus  Brain,Prostate,  function:Binds GTP, GDP and GMP.,sequence caution:Contaminating sequence. Potential poly-A sequence.,sequence caution:Unlikely isoform. Aberrant splice

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658








	Multiple transcript variants encoding different isoforms have been been found for this gene. [provided by RefSeq, Jan 2016],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



55-



## **Products Images**









Western blot analysis of lysates from HT-29 and COLO cells, using GBP3 Antibody. The lane on the right is blocked with the synthesized peptide.

## GBP3 --


nce technology Co.,Ltd

025-5229-8998 监督电话: 15950492658









Immunohistochemical analysis of paraffin-embedded human small intestinal carcinoma tissue. 1,primary Antibody was diluted at 1:200(4° overnight). 2, Sodium citrate pH 6.0 was used for antigen retrieval(>98° C,20min). 3,Secondary antibody was diluted at 1:200

Nanjing BYabscience technology Co.,Ltd