



# MARK4 Polyclonal Antibody

<b>Catalog No</b>	BYab-14826
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	IF;ELISA
<b>Gene Name</b>	MARK4
<b>Protein Name</b>	MAP/microtubule affinity-regulating kinase 4
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human MARK4. AA range:461-510
<b>Specificity</b>	MARK4 Polyclonal Antibody detects endogenous levels of MARK4 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	MARK4; KIAA1860; MARKL1; MAP/microtubule affinity-regulating kinase 4; MAP/microtubule affinity-regulating kinase-like 1
<b>Observed Band</b>	83kD
<b>Cell Pathway</b>	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytoplasm, cytoskeleton, microtubule organizing center . Cytoplasm, cytoskeleton, cilium basal body . Cytoplasm, cytoskeleton, cilium axoneme . Cytoplasm . Cell projection, dendrite . Localized at the tips of neurite-like processes in differentiated neuroblast cells. Detected in the cytoplasm and neuropil of the hippocampus. .
<b>Tissue Specificity</b>	Ubiquitous. Isoform 2 is brain-specific (PubMed:11326310). Expressed at highest levels in brain and testis. Also expressed in heart, lung, liver, muscle, kidney and spleen (PubMed:14594945).
<b>Function</b>	catalytic activity:ATP + a protein = ADP + a phosphoprotein.,similarity:Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. MARK subfamily.,similarity:Contains 1 KA1 (kinase-associated) domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 UBA domain.,tissue specificity:Ubiquitous. Isoform 2 is brain-specific.,

**Nanjing BYabscience technology Co.,Ltd**



## Background

microtubule affinity regulating kinase 4(MARK4) Homo sapiens This gene encodes a member of the microtubule affinity-regulating kinase family. These protein kinases phosphorylate microtubule-associated proteins and regulate the transition between stable and dynamic microtubules. The encoded protein is associated with the centrosome throughout mitosis and may be involved in cell cycle control. Expression of this gene is a potential marker for cancer, and the encoded protein may also play a role in Alzheimer's disease. Pseudogenes of this gene are located on both the short and long arm of chromosome 3. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Dec 2010],


## matters needing attention

Avoid repeated freezing and thawing!

## Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Immunofluorescence analysis of A549 cells, using MARK4 Antibody. The picture on the right is blocked with the synthesized peptide.

**Nanjing BYabscience technology Co.,Ltd**

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658