



Hox-A1 Polyclonal Antibody

| | |
|--------------------|---|
| Catalog No | BYab-15758 |
| Isotype | IgG |
| Reactivity | Human;Mouse;Rat |
| Applications | WB;IHC;IF;ELISA |
| Gene Name | HOXA1 |
| Protein Name | Homeobox protein Hox-A1 |
| Immunogen | The antiserum was produced against synthesized peptide derived from human HOXA1. AA range:171-220 |
| Specificity | Hox-A1 Polyclonal Antibody detects endogenous levels of Hox-A1 protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/20000.. IF 1:50-200 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | HOXA1; HOX1F; Homeobox protein Hox-A1; Homeobox protein Hox-1F |
| Observed Band | 37kD |
| Cell Pathway | Nucleus . |
| Tissue Specificity | Ovary,Skin, |
| Function | disease:Defects in HOXA1 are the cause of Athabaskan brainstem dysgenesis syndrome (ABSD) [MIM:601536]; also known as Narvajo brainstem syndrome. This syndrome is characterized by horizontal gaze palsy, sensorineural deafness, central hypoventilation, and developmental delay. Some patients had swallowing dysfunction, vocal cord paralysis, facial paresis, seizures, and cardiac outflow tract anomalies.,disease:Defects in HOXA1 are the cause of Bosley-Salih-Alorainy syndrome (BSAS) [MIM:601536]. Affected individuals show horizontal gaze abnormalities, deafness, facial weakness, vascular malformations of the internal carotid arteries and cardiac outflow trac. Some patients manifest mental retardation and autism spectrum disorder. In contrast to individuals with ABSD, central hypoventilation is not observed in individuals with |

Nanjing BYabscience technology Co.,Ltd



BSAS.,function:Sequence-specific transcription factor which is par

Background

In vertebrates, the genes encoding the class of transcription factors called homeobox genes are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosome 7 and encodes a DNA-binding transcription factor which may regulate gene expression, morphogenesis, and differentiation. The encoded protein may be involved in the placement of hindbrain segments in the proper location along the anterior-posterior axis during development. Two transcript variants encoding two different isoforms have been found for this gene, with only one of the isoforms containing the homeodomain region. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from HeLa, COLO, and 293 cells, using HOXA1 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabs science technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658